

Stratospheric Observatory for Infrared Astronomy (SOFIA)

Observing Cycle 3

Call for Proposals

May 23, 2014

Updated June 20, 2014

This document and all other information pertaining to SOFIA observing Cycle 3 may be found at http://www.sofia.usra.edu/Science/proposals/cycle3.

Key Dates

Release of Call for Proposals May 23, 2014

Call for Proposals Update on Website June 20, 2014

Proposals Due July 19, 2014 06:59 UTC

(July 18, 2014 23:59 PDT)

Anticipated Announcement of Selections Early October 2014

Cycle 3 Period 1 March 2015 – 31 January 2016









Table of Contents

Change Log:	3
. SOFIA Observing Cycle 3 Program Description	4
1.0. New policies and Capabilities for Cycle 3:	
1.1. Introduction:	4
1.2. The SOFIA Program	
1.3. Available Instruments and Observation Configurations	
1.3.1. EXES supported configurations in Cycle 3	
1.3.2. FIFI-LS supported configurations in Cycle 3	
1.3.3. FLITECAM supported configurations in Cycle 3	
1.3.4. FLITECAM/HIPO supported configurations for Cycle 3	
1.3.5. FORCAST supported configurations in Cycle 3	
1.3.6. GREAT supported configurations in Cycle 3	
1.3.7. HIPO supported configurations in Cycle 3	
1.4. Cycle 3 Schedule	
1.5. General Guidelines and Policies	17
1.5.1 Proposal Process	
1.5.2. Who May Propose	
1.5.3. Late Proposals	
1.6. Data rights and distribution.	
1.7. Targets for Observations	
1.7.1. Reserved Observations.	
1.8. Proposal Evaluation and Selection Process	
1.9. Funding for U.Sbased Investigators	
1.10. Proposer Participation in Observations	
1.11. Outreach	
1.11.1 Education and Public Outreach.	
1.11.2 Press Releases and Presentations	
1.11.3 Internal NASA Presentations	
2. Proposal Preparation and Submission	
2.1. Types of Programs	
2.1.1. Regular Programs	
2.1.2. Survey Programs	
2.1.3. Target of Opportunity Programs	
2.2. Proposal Preparation	
2.2.1 The SOFIA Proposal Tool (SPT)	
2.2.2 Proposal Text Sections (To be uploaded as a single PDF file)	
2.2.3. Exposure time estimates	
2.2.4. Sky availability during Cycle 3	
2.3. Proposal Submittal	
3. Observations and Data	
3.1. Flight Planning & Target Prioritization	
3.2. Data Processing, Calibration and Distribution	
3.2.1 Data Processing, Archiving and Distribution	29








3.2.2 Calibration.	29
4. Contacts and Further Information	30
Appendix A1 - EXES Cycle 3 Reserved Observations Catalog (ROC)	31
Appendix A2- FLITECAM Reserved Observations Catalog (ROC)	
Appendix A3 - FORCAST Imaging Reserved Observations Catalog (ROC)	33
Appendix A4 - FORCAST grism Reserved Observations Catalog (ROC)	
Appendix A5 - GREAT Cycle 3 Reserved Observations Catalog (ROC)	
Appendix A6 - FIFI-LS Reserved Observation Catalog	

Change Log:

May 23, 2014: Original release

June 20, 2014: Scheduled update release

Removed text reference to the GREAT M-channel, which is not offered in Cy 3.







1. SOFIA Observing Cycle 3 Program Description

1.0. New policies and Capabilities for Cycle 3:


- The nominal Cycle 3 observing period will be from 1 March 2015 –31 January 2016. [Section 1.4]
- The top ~5% in time of accepted programs may be carried over between cycles [Section 2.1]
- EXES and FIFI-LS have been added to the available instrument suite [Sections 1.3.1 & 1.3.2]
- FLITECAM long wavelength observations, $\lambda > 3.5 \mu m$ are offered in shared risk mode for Cycle 3 [Section 1.3.3]
- GREAT will offer the H-channel, covering the 63 µm [O I] line. [Section 1.3.6]
- Up to two Southern Hemisphere instrument series are under consideration for Cycle 3, dependent on available funding. [Section 2.2.4]
- The SOFIA Science Center allocates at least 20% of the observing time for high scientific impact proposals in excess of 40 hours. Up to 100% of the time may be assigned to these larger proposals if the Time Allocation Committee judges them sufficiently meritorious. [Section 1.1 & 1.8]
- A new "Scientific Context" section is now a required component of the proposal. [Section 2.2.2]

1.1. Introduction:

Due to the uncertainties in the NASA budget - and in particular the SOFIA project allocations - for Fiscal Year 2015 and beyond, the implementation of this Call for Proposals, and subsequent observations, is contingent on the budget appropriation for the SOFIA project for that time period. Specifically, the amount of observing time available, any southern deployments, and potentially also the number of supported instrument modes will depend on the funding levels.

The Stratospheric Observatory for Infrared Astronomy (SOFIA) is pleased to invite proposals for Cycle 3 observations, which will take place in the time period 1 March 2015 - 31 January 2016.

SOFIA cycle 3 observations will take place in a number of Science Flight Campaigns over the duration of the cycle. The campaigns will be interspersed with aircraft maintenance and instrument commissioning. Instruments that will be commissioned in 2015 are the multi-beam heterodyne receiver upGREAT and the far infrared camera and polarimeter HAWC+. These new capabilities will be offered in Cycle 4. Up to two Southern Hemisphere observing series are under consideration for the Cycle 3 time period, nominally in the summer of 2015.







For Cycle 3, the available instruments will be the mid-infrared high-resolution spectrograph EXES¹, the integral-field spectrometer FIFI-LS², the near-infrared camera FLITECAM³, including its grism modes, the mid-infrared camera FORCAST⁴, including its grism modes, the heterodyne spectrometer GREAT⁵, and the high-speed optical photometer HIPO⁶. The combination of FLITECAM/HIPO will also be available. The mid-infrared high-resolution spectrometer EXES and the integral-field spectrometer FIFI-LS are being commissioned during Cycle 2, and are offered for General Investigator observing in Cycle 3, contingent on the successful instrument commissioning.

This Call for Proposals solicits proposals for approximately 450 hours of observing time, conditional on the outcome of the US federal FY2015/2016 budgets and NASA allocations. The Universities Space Research Association (USRA) is issuing this call on behalf of NASA. Funding to support the selected applicants will also be issued through USRA. Contingent on budget confirmation and NASA approval, the total available General Investigator funding available for Cycle 3 is expected to be approximately \$1.4M.

The Call is open to all qualified astronomers, in the U.S. and outside the U.S., except for those currently affiliated with German institutions. Astronomers with a German professional affiliation must participate through a separate German Call for Proposals administered by the German SOFIA Institute (Deutsches SOFIA Institut; DSI) on behalf of the German Aerospace Center (Deutsches Zentrum für Luft und Raumfahrt; DLR). DSI personnel, even if based in the U.S., are considered affiliated with a German institution and must submit any proposals to the DLR queue. Scientists based in Germany and affiliated with the European Southern Observatory (ESO) or the European Space Agency (ESA) are considered not to be affiliated with a German institution, and may respond to this Call for Proposals.

In addition to the 450 hours available under this Call, approximately 50 hours of observing time will be available to German General Investigators through the DLR Call. An additional 7% of the Research Hours on SOFIA are set aside as Director's Discretionary Time, and the Science Instrument development teams have an allocation of

http://www.sofia.usra.edu/Science/instruments/instruments exes.html


http://www.sofia.usra.edu/Science/instruments/instruments_fifils.html

http://www.sofia.usra.edu/Science/instruments/instruments_flitecam.html

http://www.sofia.usra.edu/Science/instruments/instruments_forcast.html

 $http://www.sofia.usra.edu/Science/instruments/instruments_great.html$

http://www.sofia.usra.edu/Science/instruments/instruments_hipo.html





¹ Echelon-Cross- Echelle Spectrograph

² Field Imaging Far-Infrared Line Spectrometer

³ First Light Infrared Test Experiment CAMera,

⁴ Faint Object infraRed CAmera for the SOFIA Telescope,

⁵ German REceiver for Astronomy at Terahertz frequencies,

⁶ High-speed Imaging Photometer for Occultation,



Guaranteed Time as specified in the SOFIA Science Utilization Policies⁷. Calibration observations are part of the observatory overhead and the required time is accounted for when calculating the total observing time solicited herein.

A small number of SOFIA Science Center staff members will be directly involved in the proposal solicitation process for Cycle 3, and will therefore recuse themselves from proposing. Other SOFIA Science Center staff members may propose. Only researchers with a U.S. affiliation are eligible to receive financial support through this solicitation.

All proposals that are considered to be scientifically well justified through scientific peer review will be considered for selection. Preference will be given to substantial investigations that demonstrate significant scientific impact from SOFIA observations. Specifically, the SOFIA Science Center intends to allocate at least 20% of the available observing time for one or more proposals in excess of 40 hours observing time if they are considered scientifically compelling by the Peer Review Committee.

1.2. The SOFIA Program

SOFIA is a joint project of NASA and DLR. SOFIA Science Mission Operations (SMO), located primarily at the NASA Ames Research Center, Moffett Field, California, is responsible for the scientific operation of the observatory. The SMO is operated by USRA under contract to NASA. The Deutsches SOFIA Institut (DSI), located at the University of Stuttgart, is the primary interface between SOFIA and the German astronomical community. The SOFIA aircraft operations are managed by the NASA Neil Armstrong Flight Research Center. The aircraft itself has its home base at Building 703 of the Neil Armstrong Flight Research Center (formerly the Dryden Airborne Operations Facility, DAOF) in Palmdale, California.

SOFIA is a 2.7m telescope, with an effective, unvignetted, diameter of 2.5m, housed in a Boeing 747-SP aircraft. Observations are typically carried out at altitudes between 11.9 km (39,000 ft) and 13.7 km (45,000 ft.). These altitudes place the observatory above at least 99% and up to 99.8% of the obscuring atmospheric H_2O vapor. The observatory can operate in the 0.3-1600 μ m wavelength range, and the seven first generation instruments (six of which are offered in this Call) cover the range 0.3-250 μ m. These instruments will provide imaging and spectroscopic capabilities for a wide range of scientific investigations.

Descriptions of the instruments can be found at http://www.sofia.usra.edu/Science/instruments/.

A number of example science cases for SOFIA have been developed over the years. Design Reference Mission Case Studies may be found at http://www.sofia.usra.edu/Science/science_cases/. A number of presentations on SOFIA

⁷ Available at http://www.sofia.usra.edu/Science/documents/index.html



DLR



science can be found at the SOFIA Speakers Bureau page at http://www.sofia.usra.edu/Science/speakers/.

These science cases illustrate the breadth of potential SOFIA capabilities. Two special issues reporting observations with SOFIA performed during the Early Science phase were published by the Astrophysical Journal Letters (Vol 749, No. 2) and Astronomy and Astrophysics (Vol. 542). An overview of SOFIA is presented in Young et al. 2012, ApJ, 749, L17: "Early Science with SOFIA, the Stratospheric Observatory For Infrared Astronomy" and in Temi et al. 2014, "The SOFIA Observatory at the Start of Routine Science Operations: Mission Capabilities and Performance" (ApJ, in press).

1.3. Available Instruments and Observation Configurations

Six instruments will be available for Cycle 3 observations: EXES, FIFI-LS, FLITECAM, FORCAST, GREAT and HIPO. The FLITECAM/HIPO combination will also be offered. For the purposes of this Call for Proposals, the Focal Plane Imager (FPI+; the guide camera) is not considered a Science Instrument. While the scientific or technical justification for a program may, therefore, not be reliant on the use of data from the FPI+, the imaging data from the camera are routinely stored in the Housekeeping data stream and can be retrieved from the Science Data Archive.


There are a number of observation configurations available or planned for each of the instruments. The following sections describe the observation configurations available for Cycle 3.

Details are available in the Observers Handbook for Cycle 3 which can be found at http://www.sofia.usra.edu/Science/proposals/cycle3/.

Observations with FORCAST, FLITECAM, GREAT, EXES and FIFI-LS may be contained within the same proposal. Observations with HIPO or FLITECAM/HIPO observations must be in separate proposals.

Each of the SOFIA Science Instruments falls into one of three classes: Facility-class Science Instruments (FSI), Principal Investigator-class Science Instruments (PSI) or Special Purpose Principal Investigator-class Science Instruments (SSI). The different classes of instruments require different levels of interaction among the proposer, the science instrument team, and the SMO staff providing support and are governed by the "SOFIA Science Utilization Policies of the Stratospheric Observatory for Infrared Astronomy (SOFIA)" available at http://www.sofia.usra.edu/Science/documents/.

Facility-class Science Instrument (FSI) — A general purpose, reliable and robust instrument that provides state-of-the-art science performance at the conclusion of commissioning and upon acceptance of the instrument by the SOFIA project. FSIs will be operated and maintained by the SMO in support of General Investigators (GIs). No







direct interactions with the instrument team are required to propose for or to use the instrument.

For Cycle 3, FORCAST and FLITECAM are considered FSIs.

Principal Investigator-class Science Instrument (PSI) — A general purpose instrument that is developed and maintained by the instrument team throughout its useful operating life. PSIs will be operated by the Science Instrument team members, both for their own observations as well as for those of successful GIs. Proposers do not need to consult with the PSI Instrument team before submitting their proposals. However, GIs are encouraged to interact with the Instrument team early, since this maximizes the chances for successful observations. Guest Investigators will receive calibrated data from the EXES, FIFI-LS and GREAT teams.

For Cycle 3, EXES observations in all modes are PSI modes

Proposers are encouraged to work closely with the EXES team in the definition and execution of the observations, and to submit questions through the SOFIA Help desk (<u>sofia help@sofia.usra.edu</u>) in order to evaluate the feasibility of their Cycle 3 projects.

For Cycle 3, FIFI-LS observations are considered PSI modes.

The transition from PSI to FSI status for FIFI-LS is expected to occur during Cycle 3, but for the purpose of this call, all proposals should be generated assuming a PSI status of the instrument. For publications that involve FIFI-LS observations, the FIFI-LS PI may designate up to 3 coauthors.

For Cycle 3, all GREAT observations are considered PSI modes.


For publications that involve GREAT observations, the GREAT PI may designate up to 3 co-authors.

Special Purpose Principal Investigator-class Science Instrument (SSI) — A special purpose instrument, specifically designed for a particular type of observation not possible or practical with FSIs or PSIs. It is expected that the Science Instrument Team will operate an SSI. GIs can only use an SSI by partnering with the Instrument Team prior to proposal submission.

For Cycle 3, HIPO and the FLITECAM/HIPO combination are considered SSI.

1.3.1. EXES supported configurations in Cycle 3

Completion of EXES commissioning is planned for late in 2014, after the proposal selection for Cycle 3. Therefore, the execution of accepted EXES observation is contingent on the successful completion of these activities. Required deviations from the assumptions made here and in other supporting documents will be decided by the SMO Director and communicated to selected PIs.





EXES observations are defined by the observing modes, the spectroscopic configuration, and the central wavelength. The following EXES modes are available for Cycle 3:

Observing modes:

- 1. Nod mode
 - On-slit nod: Source moved between two points along slit for sky subtraction
 - Off-slit nod: Source moved off slit for sky subtraction
- 2. Map mode
 - Stepped maps with sky subtraction using edge of map or separate sky observation

Spectroscopic Configurations:

- 1. High-Medium
 - Echelon plus Echelle grating at angles 35-65°
- 2. High-low
 - Echelon plus Echelle grating at angles 10-25°
- 3. Medium (long-slit)
 - Echelle grating alone at angles 35-65°
- 4. Low (long-slit)
 - Echelle grating alone at angles 10-25°


Map mode is available for all spectroscopic configurations. For the HIGH_MEDIUM configuration, on-slit nodding is only available if the slit is longer than about four times the FWHM of the PSF. A more precise number will be determined during instrument commissioning. Slit lengths in this mode are a strong function of wavelength, and users must consult the online Exposure Time Calculator (ETC) to determine if on slit nodding is possible.

EXES Configuration summary (See Observers' Handbook for details)

Configurations	Wavelength (µm)	Slit	Max. Resolving Power
High-medium	4.5 - 28.3	(1.4-3.2)"x(4-40)"	100,000
High-low	4.5 - 28.3	(1.4-3.2)"x(<12)"	100,000
Medium	4.5 - 28.3	(1.4-3.2)"x(25-180)"	20,000
Low	4.5 - 28.3	(1.4-3.2)"x(25-180)"	4,000

For the high-resolution modes, there is non-continuous spectral coverage for $\lambda > 19 \mu m$, but the central wavelength can be tuned so that lines of interest do not fall in the gaps (see the SOFIA Observers' Handbook for details).

The spectroscopic observing modes have not yet been fully demonstrated in flight. For this reason, EXES observations in Cycle 3 are "shared risk." In particular, the LOW







configuration may be strongly affected by saturation effects, which has not been tested yet in flight. Therefore, the efficiency and sensitivity cannot be accurately estimated prior to this Call for Proposals. Proposers should use the information on the SOFIA website and the EXES exposure time calculator available at: http://irastro.physics.ucdavis.edu/exes/etc/

to evaluate their proposed observation parameters.

The online information will be updated as the analysis of commissioning data is ongoing. For the purposes of this Call for Proposals, the June update to this Call shall be considered definitive with regards to observing mode availability and capabilities.

1.3.2. FIFI-LS supported configurations in Cycle 3

FIFI-LS observation configurations consist of the center wavelength for each of the two Littrow spectrometers, and a selected observing mode. FIFI-LS has two independently settable spectrometers that cover the spectral ranges 50 - $125~\mu m$, and $105-200~\mu m$, respectively. The spectrometers are fed by one of two dichroics that allow simultaneous observations of the same target in the two ranges. For line selection in the wavelength overlap region (105 - $125~\mu m$) one of two available dichroic filters has to be selected, during the detailed observation set-up in Phase II. Similarly the need for spectroscopic and spatial dithering will be left for phase II inputs. For compact sources, the Map Area should be set to zero even if dithering is intended.


Observing modes:

- 1. Beam Switch mode (BSW). This is a nod-match-chop mode suitable for isolated compact sources. For such sources this is the most efficient observing mode.
- 2. Chop-offset-nod mode (C2NC2). This mode is suitable for extended sources where a symmetric chop angle would induce excessive coma in the observations.
- 3. Mapping mode (MAP). Similar to the C2NC2 mode, but with spatial offsets between the pointing centers. For mapping observations a non-zero Map Area must be chosen.

Please see the SOFIA Observers' Handbook for further details of observing modes and instrument capabilities.

1.3.3. FLITECAM supported configurations in Cycle 3

FLITECAM observation configurations consist of observing modes and filter selections in imaging mode and observing modes and grism selections in spectroscopy mode. The following are available for Cycle 3.







IMAGING

Observing modes:

- 1. Stare (Options of 5 and 9 point dither)
- 2. Nod Off Array

Filters:

Broad-band: J, H, K, L, M.

Narrow-band: Pa α , Pa α continuum, Water Ice (3.08 μ m), PAH (3.29 μ m), L_{narrow}, M_{narrow}

The Order Sorting Filters (H_W , K_W , K_{long} ; used for grism spectroscopy) are **not** offered as stand alone filters.


As the instrument commissioning was performed in the combined FLITECAM/HIPO mode, with its associated elevated long-wavelength background, the performance in L and M band (regular and narrow filters), and the LMA grism mode (grism A with the L+M order sorting filter) have not been verified. Hence these modes, as well as the PAH (3.29µm) and Water Ice (3.08µm) filters, and the "LMB" grism mode, are shared risk observations. If the elevated backgrounds are not (fully) rectified, accepted proposals requesting these filters will be executed only after a case-by-case analysis of their actual feasibility.

SPECTROSCOPY

Observing modes:

1. Nod

The spectroscopic observing mode has not yet been fully evaluated, so the efficiency and sensitivity cannot be accurately estimated prior to this Call for Proposals. Proposers should use the information on the SOFIA website to determine sensitivity and should not expect performance to exceed those numbers.







Grisms

The FLITECAM slit mask has a 2' long slit, half of which is 2" wide and half 1" wide. The spectral resolution in the 2" slit is approximately 1200.

Designation	Grism	Order	Order	Wavelength
			Sorting	Coverage
			Filter	[µm]
FLT_A1_LM	A	1	L+M	4.395 - 5.533
FLT_A3_Hw	A	3	H_{wide}	1.497 - 1.977
FLT_A2_KL	A	2	K _{long}	2.216 - 2.784
FLT_B3_J	В	3	J	1.14 - 1.424
FLT_B2_Hw	В	2	H_{wide}	1.649 - 2.076
FLT_B1_LM	В	1	L+M	3.07 - 4.16
FLT_C4_H	C	4	H	1.445 - 1.801
FLT_C3_Kw	С	3	K _{wide}	1.872 - 2.346
FLT_C2_LM	С	2	L+M	2.756 - 3.467

1.3.4. FLITECAM/HIPO supported configurations for Cycle 3

FLITECAM observation can be performed in the short wavelength modes above when co-mounted with HIPO. Due to the additional background from the warm optics in this configuration, the L, M, L_{narrow}, M_{narrow} and FLT_A1_LM modes of FLITECAM are not available when FLITECAM and HIPO are co-mounted.


1.3.5. FORCAST supported configurations in Cycle 3

FORCAST observation configurations consist of observing modes and filter selections in imaging mode and observing modes and grism selections in spectroscopy mode. The following are available for Cycle 3:

IMAGING

Observing modes:

- 1. Two position chop and nod (C2N), which may be implemented as either Nod-Match-Chop or Nod-Perpendicular-Chop
- 2. Two position large-amplitude chop (2-8 arcmin) with large nod offsets (C2NC2)



NASA =



Filters:

The full complement of filters available for the FORCAST Short Wavelength Camera (SWC; listed below) exceeds the number of available filter wheel slots. A nominal filter set has been selected for Cycle 3. Depending on the proposal requests, this nominal set may be revised prior to the start of the cycle. If required, the SOFIA Project will consider one filter swap during the duration of Cycle 3. Proposals requesting any of the non-nominal SWC filters should, in addition to justifying their filter preference, discuss the impact on the proposed science if only the "nominal" filter set is available.

For the Short Wavelength Camera (SWC) the nominal filter set for Cycle 3 is: $6.4, 6.6, 7.7, 11.1, 19.7, 25.4 \mu m$ Additional, potentially available, filters for the SWC are: $5.4, 5.6, 8.6, 11.3, 11.8 \mu m$

For the Long Wavelength Camera (LWC): 31.5, 33.6, 34.8, 37.1 µm

Dichroic:


For Cycle 3, FORCAST can be used in a single channel configuration or dual channel configuration. In dual channel configuration, a dichroic is used to split the incident light towards the short and long wavelength arrays simultaneously. Any short wavelength filter can be used at the same time as any of the long wavelength filters. However, there is significant degradation of throughput for short wavelength filters less than $11~\mu m$ and greater than 30~microns in dual-channel configuration; this information is built into the sensitivity estimator (SITE).

SPECTROSCOPY

Observing modes:

- 1. C2N (two position chop and nod).
- 2. C2NC2 (two position large-amplitude chop with large nod offsets).
- 3. SLITSCAN (A non-zero Map Area must be given)

Although the observing modes for spectroscopy are the same as for imaging, the restrictions on actual chop and nod parameters would be different. (Details may be found in the Observers Handbook.) The analysis of the commissioning data for the spectroscopic observing mode is still ongoing. Hence, the efficiencies and sensitivities for these modes are estimated for this Call for Proposals. An exposure time estimator tool is available on the Cycle 3 web page.







Grisms and Slits:

Grism	Wavelength (µm)	Slit	Resolving Power		
Long Slit Spectroscopy in the Short Wavelength Camera					
FOR_G063	4.9-8.0	2.4"x191"	200		
		4.7" x191"	100		
FOR_G111	8.4-13.7	2.4" x191"	300		
		4.7" x191"	150		
Cross Di	spersed Spectroscopy is	n the Short Wavelength	Camera		
FOR_XG063	4.9-8.0	2.4"x11.25"	1200		
FOR_XG111	8.4-13.7	2.4"x11.25"	800		
Long	Slit Spectroscopy in th	e Long Wavelength Ca	mera		
FOR_G227	17.6-27.7	2.4"x191"	140		
		4.7" x191"	70		
FOR_G329	28.7-37.1	2.4" x191"	220		
		4.7" x191"	110		

Dichroic:

For Cycle 3, all FORCAST spectroscopic observations will be done using the single channel configuration.

1.3.6. GREAT supported configurations in Cycle 3


GREAT observation configurations consist of observing modes and receiver band and backend selections. The following are available for Cycle 3:

Observing modes:

- 1. Single pointing position switching (PSW)
- 2. Single pointing beam switching (BSW), chopping with the secondary
- 3. On-the-fly mapping in PSW or BSW mode (OTFMAP PSW/BSW)
- 4. Raster mapping in PSW or BSW mode (RASTERMAP_PSW/BSW)

Receiver bands:

GREAT is a dual channel receiver where the two bands are operated simultaneously. Each band (front end) can be tuned separately. The usable instantaneous bandwidth is channel-dependent. For the L1, L2 bands the noise roll-off with intermediate frequency, intrinsic to HEB detectors, limits the usable 3-dB noise bandwidth to typically 2 GHz. Each front-end is connected to a digital FFT spectrometer providing 2.5 GHz bandwidth with 88 kHz spectral resolution (equivalent noise bandwidth).







For Cycle 3 the offered configurations will be: L1/L2, and L2/H. Depending on the results of the Call for Proposals, not all combinations may be executed. The frequency ranges of the bands are:

Band L1: 1.25 – 1.52 THz

Band L2: 1.81 - 1.91 THz

Band H: 4.7 THz

Note that not all frequencies are available within the L1 and L2 bands, and availability will depend on specific performance of the local oscillators. Potential users should consult with the instrument handbook and the GREAT instrument team.

Based on the commissioning flights in May 2014, GREAT will offer the 4.7 THz (63 $\mu m)$ H-Band. Because of the limited tuning range of the Quantum Cascade Laser local oscillator, only the velocity range -25 to +90 km/s around the 4744.77749 GHz [O I] line will be available with the standard local oscillator set-up. With the negative velocity set-up, the usable velocity range is -30 to -140 km/s. Potential proposers should contact the GREAT Principal Investigator regarding non-standard tunings.

Performance for this Call will be available on the SOFIA website.

Backends:

Fast Fourier Transform Spectrometer:

XFFTS: 2 x 2.5 GHz bandwidth with 0.088 MHz ENB.


1.3.7. HIPO supported configurations in Cycle 3

A HIPO observation configuration is the combination of a detector operation mode and optical configuration. Optical configurations, which include the choice of filter sets (in each wheel), dichroic, and reimaging optics, are provided on a per-flight basis whereas detector operation modes are selected on a per observation basis. (The FLITECAM/HIPO combination has the same modes as HIPO, with the additional FLITECAM filters and grisms available.) The following are available in Cycle 3:

Observing modes:

- 1. Single frames
- 2. Basic Occultations
- 3. Fast Occultations

Filters:







U, B, V, R, I (Johnson) u', g', r', I', z' (Sloan Digital Sky Survey) Methane (890 nm)

Up to eight filters may be accommodated per filter wheel. For the dual-channel imaging mode, all combinations of filters and dichroics with overlapping transmission are available. For the red-channel (see below), all filters are available except U and u'. Filter wheel positions can also be left OPEN on any wheel if desired. All permitted filter/dichroic combinations can be used with any one of the detector operation modes for an observation.

Dichroics:

There are two dichroics available with cut-offs at 575 and 675 nm. The choice of dichroic is determined by a preference for where to split the wavelength coverage of the 'blue' and 'red' channels. For example, the 675 nm dichroic permits simultaneous observation in R and I bands in the 'blue' and 'red' CCD channels, respectively. HIPO also supports a 'red' channel only configuration (no dichroic) for higher optical throughput if two-color photometry is not required.

Custom Optical Configurations and Detector Operation Modes:

HIPO can accommodate custom filters and dichroics for specific observations. A 'bare CCD' configuration without re-imaging optics and with one filter wheel is also available. Additional engineering readout modes are also available for special purpose observations in collaboration with the HIPO instrument team.

1.4. Cycle 3 Schedule

The nominal schedule for the Cycle 3 observing program is as follows:

23May 2014 Release of Call for Proposals

20 June 2014 Call for Proposals update

18 July 2014, 23:59 PDT Proposal Submission deadline
19 July 2014, 06:59 UTC Proposal Submission deadline
Early October 2014 Proposal Selections Announced.

1 March 2015 – 31 January 2016 Cycle 3 observing period.







SOFIA observations in Cycle 3 will be conducted in a number of Science Flight Campaigns⁸ covering the periods March, 2015 - February, 2016. The detailed layout of Science Flight Series within each Science Flight Campaigns will depend on the selected proposals and instrument availability.

1.5. General Guidelines and Policies

At the time of writing of this Call for Proposals, only FLITECAM, FORCAST, GREAT, HIPO, and FIFI-LS have completed their commissioning observations, and only FORCAST imaging mode has passed its formal acceptance review. A significant part of the analysis of the commissioning data for FLITECAM, FORCAST grism-mode, and FIFI-LS still remains to be completed. Hence observations with EXES, FIFI-LS and grism mode observations with FLITECAM and FORCAST carry a larger uncertainty and may require modifications or reconsideration by the SMO Director once the commissioning data have been fully analyzed.


Observations lost due to observatory or instrument hardware or software failures, weather, or other reasons, will not automatically be rescheduled. In general, with the exception noted in Sec. 2.1, such observations will need to be re-proposed for in a subsequent Cycle.

1.5.1 Proposal Process

The SOFIA Cycle 3 proposal process will consist of two steps – Phase I and Phase II. Phase I requires the preparation and submission of a science justification, a feasibility analysis for the proposed program, and a high level description of the proposed targets and observations. This Phase I proposal will form the basis of the peer review and proposal selection by the SMO Director. Proposals that are awarded observing time based on the evaluation process described in Section 1.9 will subsequently be required to submit Phase II observation specifications following guidelines provided by the SMO. These submissions will provide the SMO and instrument PIs with the detailed definition of each observation to be executed for the program. In addition, proposers affiliated with U.S. institutions will be invited to submit a budget, based on funding guidelines provided by the SMO.

a) Science Flights - individual flights primarily devoted to obtaining astronomical science data.

- c) Science Flight Campaigns One or more science flight series, beginning and ending with a non-science, engineering activity.
- d) *Science Observing Cycles* One or more of flight campaigns that are covered by a single science Call for Proposals.





⁸ SOFIA science observing definitions:

b) *Science Flight Series* - Contiguous series of science flights, all with the same instrument.



SOFIA Cycle 3 Phase I proposals must be prepared and submitted using the SOFIA Proposal Tool (SPT) (http://dcs.sofia.usra.edu/proposalDevelopment/installSPT), which is a Java based application. The prospective proposer should download SPT to a local computer. The proposal consists of formatted information filled in via the SPT form fields (such as proposer information, scientific category, instrument, target and exposure information) and a file containing the scientific justification and other information (details in section 2.2.1), to be uploaded in pdf format. SPT is available for most commonly used platforms, including Mac OS X, Windows and Linux (http://dcs.sofia.usra.edu/proposalDevelopment/installSPT/index.jsp).

An outline of the proposal preparation process may be found in section 2 and further details about SPT in the Observers Handbook.

1.5.2. Who May Propose


Participation in the U.S. SOFIA Cycle 3 Program is open to scientists from all categories of U.S. and non-U.S. organizations, including educational institutions, industry, nonprofit institutions, NASA Centers, and other Government agencies. Members of German organizations, including DSI staff stationed in the U.S., must participate through the DSI led program.

Each SOFIA Cycle 3 proposal must identify a single Principal Investigator (PI) who assumes responsibility for the conduct of the scientific investigation. Proposal Co-Investigators must have well defined roles in the investigation, which will be evaluated as part of the proposal review process. Following selection by the SMO Director, the SMO staff will communicate formally only with the PI (or a person designated by the PI) of each proposal, except for funding issues where communications will be primarily with the institutional Sponsored Research Office (or equivalent). It is the responsibility of the PI (or designee) to provide the SOFIA project, in a timely manner, all information necessary for implementing observations, and to respond to any questions concerning observational constraints or configurations.

Note: those with a German professional affiliation may participate as Co-Investigators on proposals submitted in response to this Call. They may not be Principal Investigators on the proposals, nor in any way be designated as the contact or lead investigator.

1.5.3. Late Proposals

Consistent with USRA and NASA policy, no late proposals will be considered. A proposal will be considered "on time" only if all necessary components have been received by the published deadline. Note that processing delays at the proposer's home institution, shipment delays of the proposal, or Internet delays, do not excuse the late submission of a proposal.







1.6. Data rights and distribution

All scientific data from SOFIA observations will be distributed via the SOFIA Data Cycle System's Science Archive

(https://dcs.sofia.usra.edu/dataRetrieval/SearchScienceArchiveInfo.jsp)

All data will be archived as Level 1 data (raw). Where appropriate, Level 2 (corrected for instrumental and atmospheric effects) and Level 3 (flux calibrated) will also be archived.

These data will be accessible to the general community after a proprietary period of twelve months, starting at the ingestion of the calibrated data into the archive.

1.7. Targets for Observations

All scientifically valid observations may be proposed for, with the exception of those duplicating the ones in the "Reserved Observations Catalog" lists (Appendix A), designated by each Science Instrument team. Duplication of observations obtained in earlier cycles are generally not allowed, and if proposed for, must be identified as such, and the necessity for duplication must be explicitly justified.


Observations that are part of an active Cycle 2 SOFIA program, but which have not been executed at the time of the Cycle 3 proposal deadline may be re-proposed, but must be clearly identified as such. If the Cycle 2 observation is executed and the Cycle 3 request qualifies as a duplication (target, mode and exposure time) the latter will be disallowed. The proposal may provide potential substitute targets, but must then clearly discuss the relative scientific justification for the original target[s] and the substitute targets.

Non-sidereal tracking of SOFIA observations has been implemented, but only limited operational experience has yet been gained. Such observations therefore carry a larger risk. Proposals of such observations are allowed, but the SMO Director will decide in each individual case whether an accepted observation is to be executed.

1.7.1. Reserved Observations

As part of the instrument development contract, the instrument teams were awarded a limited amount of Guaranteed Time Observing (GTO). To protect the interests of the instrument development teams, the SOFIA project has allowed the Cycle 3 instrument teams to set aside a limited number of targets and associated exposure times as "Reserved Observations". These reserved observations may not be proposed for in Cycle 3.

A Reserved Observation consists of the combination of position on the sky, instrument, observation configuration, and length of observation. The observation configuration encompasses the basic scientific intent of the observation by specifying, for example, the wavelength range for broad-band photometry or grism spectroscopy or the frequency of observation for GREAT. Similarly, proposed observations are considered duplicate to







previously observed observations if they duplicate the combination of position on the sky, instrument, observation configuration, and length of observation. Hence, observations of the same target but in different filters are not considered duplicates.

The Reserved Observation Catalogs for the instruments are independent of each other. The Reserved Observation list only applies to Cycle 3, and the Instrument PIs will have the opportunity to revise their Reserved Observation Catalog prior to subsequent proposal calls.

Requests for an observation in the Reserved Observation Catalogs may only be proposed for scientific investigations that cannot be accomplished, or were not intended to be accomplished, by the Instrument Team. The spirit of these restrictions is to protect science objectives, and therefore observations that closely resemble any in the Reserved Observation Catalogs would also be disallowed.

If a reserved observation is proposed for, the justification for such a duplication must be clearly addressed in the proposal. At a minimum, any such proposals must aim to achieve a signal-to-noise ratio of twice that expected from the Reserved Observation or have a scientifically justified duplication such as for temporal variability studies. (Since the Reserved Observation Catalogs for the instruments are independent of each other, a target in one Catalog may be proposed for using a different instrument, without the above exposure time minimum.) Final determination of acceptability of proposed observations rests with the SMO Director.

The Reserved Observations Catalogs can be found in Appendix A.


1.8. Proposal Evaluation and Selection Process

Proposals submitted in response to this Call will be evaluated in a competitive peer review. The peer review panel, including its chair, will be recruited from the astronomical community and be subject to the standard NASA procedures and rules.

The following factors will be used in evaluating proposals for the SOFIA Cycle 3 Program.

- 1. The overall scientific merit of the proposed investigation.
- 2. The broader scientific impact of the investigations to astronomy.
- 3. The feasibility of accomplishing the objectives of the investigation.
- 4. The degree to which the investigation uses SOFIA's unique capabilities.
- 5. The competence and relevant experience of the Principal Investigator and any collaborators to carry the investigation to a successful conclusion.

The scientific review panels will be given an assessment of the technical feasibility of each proposal as determined by the SMO. After acceptance of an observing program, successful proposers must provide the required inputs to detailed observing plans for





submission to the SMO. Instructions for completing these Phase II inputs will be distributed to the PIs of the selected proposals.

Proposals of all sizes will be considered. For Cycle 3, the SMO encourages substantial investigations with significant impact and plans to allocate at least 20% of the available observing time to high scientific impact proposals in excess of 40 hours. Up to 100% of the time may be assigned to these larger proposals if the Time Allocation Committee judges them sufficiently meritorious.

The SOFIA project reserves the right to select only a portion of a proposer's investigation, in which case the PI of the proposal will be given the opportunity to accept or decline the implementation of the partial selection.


Because of the complicated process of flight scheduling involving sky visibility, instrument availability, and the need to produce efficient flight plans, selection of an investigation does not guarantee observation. At the discretion of the SMO director, an appropriate over-subscription of the available flight times may be accepted via the peer review process with an associated prioritization, which will allow for contingencies in flight planning. The SMO Director will approve the implementation of the observing prioritization and target selection.

1.9. Funding for U.S.-based Investigators

Limited funds for awards under this solicitation are expected to be available to investigators at U.S. institutions subject to the annual NASA budget cycle. Successful proposers at U.S. institutions, including U.S. Co-Investigators on successful non-U.S. proposals, will be eligible for funding.

Budgets should not be submitted with the proposals in response to this Call. The selected investigators will receive a funding guideline from the SOFIA Science Center based on the scope of the approved observing program and the available budget for the SOFIA Cycle 3 program. A budget summary and narrative description of how these funds will be used must be submitted after the receipt of the guideline. The deadline for budget submittal will be announced after the proposal selection and included in the funding guidelines document. An institutional signature will be required when a budget is submitted.

Grants disbursement will be implemented in two steps. A first installment of ~\$2k will be disbursed at the time of proposal selection to support the work required to generate the detailed observing plan in Phase II. The remaining part of the grant will be disbursed at the time of the initial observations in the program. Deviations from this disbursement schedule can, for exceptional circumstances, be requested through the SMO Director.







1.10. Proposer Participation in Observations

SOFIA GIs are encouraged to participate in the on-board observations and can contribute to the success of their observations. However, observations from many different programs are usually executed on any given flight. This has several impacts on GI inflight participation: 1) Only a limited number of observations in the GIs program are likely to be executed on any given flight. 2) While many optimizations of a given observation are possible in-flight, the ability to interactively modify a program is limited to the specific observation. Changes that would affect the remainder of the flight plan (e.g. target changes), or that could cause conflicts with other accepted programs (such as filter settings not originally awarded to the current program), will generally not be allowed. 3) With the many different required and requested flight crew complements the number of "Astronomer seats" on any given flight is limited. For a given flight, if the number of GIs requesting seats exceeds the number available, then the SMO Director will decide on which GIs will be invited on that flight.


1.11. Outreach

1.11.1 Education and Public Outreach

The SOFIA project strongly encourages observing programs to include an Education & Public Outreach (E&PO) component, supported by the SOFIA Outreach office. The proposer does not need to add any E&PO text to the submitted observing proposal. For selected programs, the SOFIA Outreach staff will contact the program PI during the Phase II stage, to collaborate in designing an E&PO program related to the program's science.

For E&PO planning purposes, proposers are requested to indicate, in their Phase I submittals, whether or not they are interested in one particular E&PO option, namely the Airborne Astronomy Ambassadors (AAA) program. Participation in the AAA program will involve a partnership between the GI team and a team of educators who will be put in contact with the scientists before the observing flight(s) of the program in question. The educator team will be trained by SOFIA Outreach to understand the planned investigation, will fly on SOFIA with a flight facilitator/escort, and then continue to partner with the GI team in a mutually agreeable fashion after the flight(s). The selection of the AAA educators is carried out through a proposal process which runs in parallel to the Cycle 3 observing proposal process.

Please note that by choosing the AAA option the proposer is not required to recruit the educators, give them materials, or fund their activities; the SOFIA Outreach office will provide all such support. If the AAA program option is chosen, and the GI chooses to participate in the SOFIA observing flight(s) for their program, the AAA team (plus escort) will accompany them on the flight(s).







The proposal cover page in SPT includes a button allowing the proposer to indicate an interest in participating in the AAA program. The SOFIA project requests that you indicate whether you are interested in participating in the AAA program in the Phase I stage since this will provide important information for the planning, execution and review of the AAA educator proposal and selection process. The choice of whether or not to indicate an interest in participating in the AAA program will not affect the evaluation of the Cycle 3 observing proposal.


1.11.2 Press Releases and Presentations

SOFIA is already capturing the imagination and attention of the press and the public. To continue this successful publicity, SOFIA observers have a responsibility to share potentially newsworthy results with the public. NASA/DLR have an interest in helping them reach a larger audience and gain a bigger impact. Specifically, NASA and DLR retain the right to be the initial organizations to issue press releases regarding SOFIA results. Therefore, if a GI believes that there is a possibility that his/her new results could be of interest to a wide public audience, they should contact the SOFIA Public Affairs officer Nicholas Veronico (nveronico@sofia.usra.edu) who will evaluate the news value of the results, communicate with NASA and DLR Headquarters, and then work with the GIs on the most suitable course of action. Releasing results without coordinating with the program or agency will prevent the result from being included in a subsequent NASA/DLR press release.

NASA and the DLR will jointly issue press releases associated with SOFIA observations during Cycle 3. The U.S. SOFIA Public Affairs officer and the German/DSI Public Affairs counterpart will coordinate the press release process. Other relevant press releases by participating organizations (including PI institutions) should be coordinated with the SOFIA program, NASA and DLR. Other presentation material based on the Cycle 3 observations can be generated by any member of the proposal team and will be considered part of the team's collective set of material. Any member of the team may use these materials (e.g., in public science talks or conference proceedings).

1.11.3 Internal NASA Presentations

Noteworthy SOFIA results are of great interest to NASA. GIs are encouraged to support internal presentations to NASA management, with the understanding that results will be made public only with the agreement of the GI. GIs will also be encouraged to make early results available in more public venues like the SOFIA website and presentations.







2. Proposal Preparation and Submission

2.1. Types of Programs

Three types of programs are solicited in response to this Call: Regular Programs, Survey Programs, and Target of Opportunity Programs. A single proposal may not mix different program types.

Starting in Cycle 3 a limited fraction (approximately 5% of allocated observing time) of the most highly ranked Regular Programs will be assigned status as "guaranteed observations" and may be carried over between cycles if they fail to be scheduled during the Cycle 3 period. Proposers do not need to, and cannot, request this status, as it will be assigned by the SMO Director as part of the proposal selection process.

With the exception described above, proposals are active only for the duration of the Cycle 3 observing period. Accepted observations not executed during Cycle 3, for whatever reason, will not be carried over to future cycles.

In addition, the SOFIA program accepts proposals for Director's Discretionary Time programs. This category is intended for short, urgent observations that could not have been foreseen at the time of the proposal call and that cannot wait for the next proposal cycle. In exceptional cases, proof-of-concept observations may be requested through the DDT path. However, a strong justification for not proposing such observations through the regular proposal process will be required. DDT proposals are not solicited through this Call for Proposals, and should be directly addressed to the SMO Director, Dr. Erick Young. Further information about the DDT program can be found at the SOFIA web site under: http://www.sofia.usra.edu/Science/proposals/DDT/instructions.html


2.1.1. Regular Programs

Observations of specific targets with known positions and timing constraints (including targets with no constraints) will constitute regular observing programs. This includes time critical observations and observations of known Solar System objects.

The intent is to execute all the observations accepted in a regular program.

2.1.2. Survey Programs

Survey proposals are intended to allow studies of a target class, as well as provide the SOFIA project flexibility in flight planning. These programs should identify a sample of targets and observations with a common scientific justification. The selection of survey proposals will be primarily judged on scientific merit, but samples with uniform sky distributions will be prioritized as they provide the best flexibility in flight planning.







The intent is that a useful fraction of the targets in a given survey program will be observed, but with no given target observation guaranteed to be executed. The proposal should discuss and justify a minimally useful fraction of completion.

2.1.3. Target of Opportunity Programs

Target of Opportunity (ToO) proposals are invited. Both programs with known targets, but unknown timing of the observations, such as observations of a specific target at an unknown time (e.g. an identified recurrent nova in outburst), and programs targeting a class of astronomical events, but with unknown targets and timings (such as observations of an as yet unidentified comet or supernova), will be considered. For ToO observations, the proposal should contain a discussion of the triggering criteria, the required turnaround time between triggering and observation, and any other timing constraints.

Since SOFIA can only observe with a single instrument at the time (with the exception of the co-mounting of HIPO and FLITECAM), rapid turn-around ToO requests with a specific instrument may be difficult to implement. Hence, ToO proposals should also address the viability and utility of observing the event/target with each of the available SOFIA instruments.

The SMO Director will have ultimate authority in recommending or rejecting the request that a selected ToO program be activated.

Observations of specific Solar System targets or events whose times of occurrence can be predicted in advance (e.g. occultations) **do not** constitute ToO observations and should not be flagged as such.


2.2. Proposal Preparation

Each Cycle 3 proposal must be prepared using the SOFIA Proposal Tool (SPT). The proposal information is entered directly, while text sections including the scientific justification and feasibility analysis should be in PDF files, uploaded via SPT⁹.

Proposals must be written in English. The length of each section of the proposal should not exceed the page limits indicated in Section 2.2.2, using single-spaced 8.5x11 inch or A4 format with 1 inch (2.5 cm) margins. Proposals must be printed to the PDF files with a font size no smaller than 11 points (about 6 characters per cm). Reviewers will only be provided the portion of each proposal that complies with the page limits.

The abstract provided using the Proposal Information form is limited to 300 words (see Section 2.1.2).

⁹ http://dcs.sofia.usra.edu/proposalDevelopment/installSPT/index.jsp



NASA = DLR



2.2.1 The SOFIA Proposal Tool (SPT)

The SOFIA Proposal Tool (SPT) provides the user with a simple form-based interface for preparing a proposal and for electronic submission to the SOFIA Science Mission Operations. After downloading the appropriate package and following the installation instructions, the user starts a new proposal by launching the SPT application. The proposer then fills out the necessary form fields including proposer information, abstract, instrument(s), and target lists. The Science and Technical Justification may be prepared using any text editor (e.g. MS Word, LaTeX, etc...) and saved as a PDF file. Using SPT, the proposer then identifies this PDF file on a local disk for attachment to the proposal summary information. When the proposal is complete, the user submits the complete proposal directly to the SMO using SPT. Upon successful submission, a unique identifier is returned for later reference. Proposals that have been submitted to the SMO can be *resubmitted* using SPT at any time up to the proposal deadline (note that old versions are not retained). On-line help for SPT is available as a pop-up function in the application.

Please note that "Blackout Periods", and "Long Term Observations" are not supported in Cycle 3 and will be ignored in the SPT form.

2.2.2 Proposal Text Sections (To be uploaded as a single PDF file)

Proposal Sections – The uploaded PDF file must contain the following sections in the order indicated for each proposed observing program. The page length limits are indicated.

1. **Scientific Context (up to 0.5 pages)** – Briefly summarize the proposed investigation with the following elements:


Context – What is the context and significance of this proposal to the broader field of astronomy?

Aims – How will the observations address the specific scientific questions in this proposal?

Methods – What are the key measurement techniques utilized in this investigation?

Anticipated results – What are the expected data sets that will be produced in this investigation?

- 2. **Scientific Justification (up to 3 pages)** Describe the scientific objectives of the proposed investigation, clearly stating the goals and their significance to astronomy, and why SOFIA data are essential to the investigation. The results and status of previous, related, SOFIA observations should be summarized. The page limit includes all text, figures and tables. An additional page for references only is allowed.
- 3. **Feasibility (up to 2 pages)** This section forms the basis for assessment of the technical feasibility of the proposed observations. The requested exposure time for each observation must be justified. The section should include the expected







target fluxes and the signal-to-noise ratio required for each observation. The source (or method) for the flux estimates, and their accuracies should be mentioned. Where applicable the spectral resolution required must be explicitly stated. Any other information about the proposed observations that would help the reviewer relate the technical needs to the scientific goals should be included in this section. Observing overheads and other indirect time estimates should follow the instructions given in the Observers Handbook. This section should also contain the justification for special calibration procedures, if they have been requested (Section 3.2.2).

4. Principal Investigator and Co-Investigator Biographical and Publication Data (one page for the PI with one additional page for CoIs). A short biographical sketch for the PI should be provided and include a list of the most recent refereed publications relevant to the scientific proposal. Short biographical data, including their roles in the proposed project, should be provided for the CoIs

2.2.3. Exposure time estimates

Estimates of exposure times for imaging with FLITECAM and FORCAST can be made using the SOFIA Instrument Time Estimator (SITE)¹⁰, a web-based tool that provides total integration time or S/N for a given instrument, filter(s), source type (point, extended, emission line) and water vapor overburden. Algorithms and assumptions used are given in the help link on the SITE webpage.

Stand-alone Exposure Time Calculator (ETC) tools for the FLITECAM- and FORCAST-grism modes and for GREAT observations have been added to the SOFIA web site under Cycle 3 information. A stand-alone exposure time estimator for EXES is available at the instrument team's web site at: http://irastro.physics.ucdavis.edu/exes/etc/

Instructions for exposure time calculations and baseline sensitivities for FIFI-LS can be found on the SOFIA web site at:

http://www.sofia.usra.edu/Science/proposals/cycle3/phase1.html

Because this call for proposals is being issued prior to in-flight commissioning of some instrument modes, the sensitivities of some of the instruments are somewhat uncertain.

2.2.4. Sky availability during Cycle 3

The sky availability for SOFIA observations is constrained by several factors, including the need to return to the Palmdale, California home base at the end of a flight and the avoidance of restricted airspace. Due to these constraints, the southernmost declination available on flights departing and landing in Palmdale is -36°. This limit is calculated based on limits of flight plans and telescope pointing. Note that, depending on the

http://dcs.sofia.usra.edu/proposalDevelopment/SITE



NASA = DLR



sensitivity of a given observation to atmospheric opacities, the limiting Declination may be significantly more stringent in practice.

Current flight rules prohibit observations when the sun is above the local horizon and require that the aircraft touch down at least 15 minutes prior to sunrise.

The instantaneous pointing of the telescope, relative to the aircraft, is restricted to $\pm 3^{\circ}$ cross-elevation (on the port (left hand) side of the plane) and elevations between 20 and 60 degrees (unvignetted). A somewhat larger vignetted elevation range is possible, but the proposer should contact the SMO regarding possible performance issues.

The SOFIA Program is evaluating the possibility of up to two Southern Hemisphere flight series sometime during the June-August 2015 period out of Christchurch New Zealand. The number of supportable flights as well as the number of instruments (one or two) available during the proposed deployments will be dependent on the funding for SOFIA during the Cycle 3 period. The over-all scientific justification (number and urgency of accepted proposals) for the different offered instruments will be a key determinant in which instruments will be available on these deployments.


Cycle 3 proposers are not expected to lay out flight plans or perform detailed visibility analysis for their proposals. Such considerations are also not needed for the phase I proposals. Flight planning will be done by the SMO staff and the instrument teams. General target availability for a specific set of dates can be judged using the SOFIA Visibility Tool (VT). The stand-alone version is available at

https://dcs.sofia.usra.edu/observationPlanning/installVT/

2.3. Proposal Submittal

Proposals must be submitted using the SPT application. Upon successful upload, the system will generate an automatic message acknowledging the submittal. A confirmation email will be sent to the address provided in the proposal. Details about the SPT may be found in the Observers Handbook.

Proposals can be resubmitted at any time before the proposal due date. Note that a resubmitted proposal **replaces** all previously submitted versions - the SMO keeps only the latest proposal submission associated with a given proposal number.



NASA = DLR



3. Observations and Data

3.1. Flight Planning & Target Prioritization

Flight planning is not part of the proposal process. However, source selection with the constraints of an airborne observatory in mind can increase the ability of a program to be scheduled. It may be expected that certain regions of the sky, such as the inner Galactic Plane, and Orion will be oversubscribed. These targets force SOFIA to fly westwards since they are towards the south (and the telescope looks out on the port - left-hand - side of the aircraft). Since, in regular operations, SOFIA has to return to Palmdale at the end of each flight, targets in the northern half of the sky will be required for roughly the same amount of time as these southern regions. (Note that southern and northern half of the sky here does not refer to south and north of the celestial equator but to sources culminating south or north of the zenith.)

The ranking by the peer review panel and selection by the SMO Director will result in a prioritized target pool, which will be provided to the SMO staff. The SMO will then produce flight plans in consultation with the instrument teams and proposal PIs. The effort will be carried out under the scientific direction of the SMO Director.

3.2. Data Processing, Calibration and Distribution

3.2.1 Data Processing, Archiving and Distribution


The SMO will be responsible for the processing of data obtained by Facility Class Science Instruments. The instrument development teams will be responsible for the data reduction for Principal Investigator Class and Special Purpose Science Instruments. All scientifically meaningful data obtained during the Cycle 3 phase will be made available to observers via the SOFIA Science Archive

For HIPO, only level 1 data will be stored at the SOFIA Science Archive. (No processing is done, so as to maintain highest fidelity for time-variability studies.)

3.2.2 Calibration

It is expected that a photometric calibration accuracy of at least 20% will be achieved for FLITECAM and FORCAST imaging and for GREAT observations, except in spectral regions of strong telluric interference. For FIFI-LS a photometric calibration accuracy of 30% is expected to be achievable, contingent on the availability of suitable calibration data.

The EXES temperature-controlled blackbody source is expected to provide flux calibration to better than 20%. However, since EXES will be commissioned at the end of 2014, the true performance of EXES is currently uncertain. Experiments focused on line profile information and those that can normalize the continuum level, or use past







observations for setting the continuum, are likely to be lower risk. Correction for the impact of Earth's atmosphere will also be done using the black body source. Corrections, using the blackbody, are expected to be accurate in wavelength regions where the atmospheric transmission is >0.50 and is spectrally smooth over the region of interest. Projects requesting additional telluric calibration will need to identify a suitable target and include the observation time required to reach the desired signal-to-noise in their proposal.

Absolute calibration for HIPO is not expected to be an issue for its scientific usage, but may be brought up by the proposer during discussions with the Instrument PI before preparing and submitting a proposal.

Cycle 3 proposals do not need to include time for calibration target observations. Observations of astronomical calibrators will be performed by the Observatory and will be allocated as general overhead and will not be counted against the awarded General Investigator time. Proposers wishing to implement specific calibration strategies may propose to do so, but must identify the specific calibration target observations to accomplish these goals and explicitly request the observing time to accomplish these observations. The calibration strategies and targets will be evaluated in the technical and science reviews, and if recommended by the review process will be treated as part of the proposal.

4. Contacts and Further Information

For further information about the Cycle 3 Call for Proposal or help in preparing proposals, please see the "Information for Researchers" (http://www.sofia.usra.edu/Science/) section of the SOFIA web site, or contact the SOFIA help desk at sofia help@sofia.usra.edu.

Questions about the SOFIA General Investigator program can be directed to the SOFIA Science Operations Manager, Dr. B-G Andersson (bg@sofia.usra.edu), or SOFIA User Support Scientist Dr. Ravi Sankrit (rsankrit@sofia.usra.edu).

For further information about the SOFIA Science project, please contact the above, or the Science Mission Operations Director, Dr. Erick T. Young (eyoung@sofia.usra.edu)







Appendix A1 - EXES Cycle 3 Reserved Observations Catalog (ROC)

Target	R.A.	Dec	$\lambda_{central}$	Config.	t _{exp}
	(2000)	(2000)	[µm]		5.1,6
IRC+10216	09:47:57.41	+13:16:43.56	5.85	High-Low	600
IRC+10216	09:47:57.41	+13:16:43.56	6.00	High-Low	600
IRC+10216	09:47:57.41	+13:16:43.56	6.15	High-Low	600
Sirius	06:45:08.92	-16:42:58.02	5.85	High-Low	900
Sirius	06:45:08.92	-16:42:58.02	6.00	High-Low	900
Sirius	06:45:08.92	-16:42:58.02	6.15	High-Low	900
Mon R2 IRS3	06:07:48.40	-06:22:55	5.705	High-Med	1600
Sirius	06:45:08.92	-16:42:58.02	5.705	High-Med	600
Mon R2 IRS3	06:07:48.40	-06:22:55	6.175	High-Med	1600
W3 IRS5	02:25:40.4	+62:05:52	5.705	High-Med	900
NGC 7538 IRS1	23:13:45.4	+61:28:09	5.705	High-Med	3600
W3 IRS5	02:25:40.4	+62:05:52	6.175	High-Med	900
NGC 7538 IRS1	23:13:45.4	+61:28:09	6.175	High-Med	3600

All observations to be taken in the "nod" Observing Mode The High-Low







Appendix A2- FLITECAM Reserved Observations Catalog (ROC)

Table 1: Target List for PAH observations

Target	RA (J2000)	Dec (J2000)	Itime + O/H ¹	Type
HB 5	17:47:56.19	-29:59:41.9	1 hr	PN
NGC 6790	19:22:56.97	+01:30:46.5	1 hr	PN
BD +30 3639	19:34:45.23	+30:30:58.9	1 hr	PN
NGC 6886	20:12:42.81	+19:59:22.7	1 hr	PN
M 3-35	20:21:03.77	+32:29:23.9	1 hr	PN
SH2-106	20:27:27.1	+37:22:39.0	1 hr	HII Region
NGC 7027	21:07:01.59	+42:14:10.2	3 hr	PN
IRAS 21282+5050	21:29:58.42	+51:03:59.8	1 hr	PN
IC 5117	21:32:31.03	+44:35:48.5	1 hr	PN
CRL 618	04:42:53.67	+36:06:53.2	1 hr	PN
Orion Bar	05:32:55.3	-05:26:50.5	4 hr	HII Region
HD44179	06:19:58.22	-10:38:14.7	1 hr	Red Rectangle
Alpha Vul	19:28:42.32	+24:39:53.6	0.5 hr	Calibrator
75 Cyg	21:40:11.11	+43:16:25.8	0.5 hr	Calibrator
HD 10074	01:39:07.73	+36:32:36.1	0.5 hr	Calibrator
HD 13746	02:14:42.16	+30:23:41.2	0.5 hr	Calibrator

^{1.} Total on-target integration time plus 30% overhead; total of 19 hours. Each observation requires a minimum of two narrow band images (on/off the PAH feature) and a 3.3 micron grism-mode spectrum. Up to ten "map" positions will be needed on the largest objects.

Table 2: Brown Dwarfs


Target	RA (J2000)	Dec (J2000)	Itime + O/H ¹	Type
Tdwarf1	07:22:27.27	-05:40:29.9	3 hr	J=16.5, W2=12.2
Tdwarf2	17:41:24.26	+25:53:19.5	3 hr	J=16.5, W2=12.3

^{1.} Total on-target integration time plus 30% overhead; total of 6 hours. Each observation requires a minimum of three 3-5 micron grism spectra (L&M plus grisms A, B, and C using ABBA nodding) as well as two broad band images (5 point dither in L and M bands).

Table 3: Paschen Alpha Imaging

Target	RA (J2000)	Dec (J2000)	Itime + O/H ¹	Type
W51	09:23:27.61	+14:30:14.77	4 hr	Star Forming Region

^{1.} Total on-target integration time plus 30% overhead; total of 4 hours. This observation requires a minimum of two narrow band images (5 point dithers in Paschen Alpha and Paschen Alpha Continuum) in at least six "map" positions. This observation is for FLIPO mode only.








Appendix A3 - FORCAST Imaging Reserved Observations Catalog (ROC)

Object Name	RA	Dec	Extent	SWC	LWC	Observing
	(J2000)	(J2000)	(arcmin)	Filter	Filter	time
				[µm]	[µm]	[hours]
Alpha Lyr	18:36:56.34	+37.1:47:01.3	3.2x3.2	0	34.8	15.83
Fomalhaut	22:57:39.05	-29:37:20.1	3.2x3.2	0	34.8	1.36
HD 141569	15:47:20.2	-03:46:12	3.2x3.2	6.3	0	0.25
HD 141569	15:47:20.2	-03:46:12	3.2x3.2	7.7	0	0.25
HD 141569	15:47:20.2	-03:46:12	3.2x3.2	8.6	0	0.25
HD 141569	15:47:20.2	-03:46:12	3.2x3.2	11.1	37.1	0.25
HD 141569	15:47:20.2	-03:46:12	3.2x3.2	11.3	37.1	0.25
Sag A West, Arches, filaments	17:45:50.5	-28:49:28	12.8x9.6	6.3	0	0.56
Sag A West, Arches, filaments	17:45:50.5	-28:49:28	12.8x9.6	6.6	0	0.56
Sag A West, Arches, filaments	17:45:50.5	-28:49:28	12.8x9.6	7.7	0	0.50
Sag A West, Arches, filaments	17:45:50.5	-28:49:28	12.8x9.6	19.7	31.4	0.35
Sag A West, Arches, filaments	17:45:50.5	-28:49:28	12.8x9.6	24.2	37.1	1.28
LBV G0.120 - 0.048	17 46 05.63	-28 51 31.9	3.2x3.2	6.3	0	0.047
LBV G0.120 - 0.048	17 46 05.63	-28 51 31.9	3.2x3.2	6.6	0	0.047
LBV G0.120 - 0.048	17 46 05.63	-28 51 31.9	3.2x3.2	7.7	0	0.042
LBV G0.120 - 0.048	17 46 05.63	-28 51 31.9	3.2x3.2	19.7	31.4	0.0083
LBV G0.120 - 0.048	17 46 05.63	-28 51 31.9	3.2x3.2	24.2	37.1	0.03
"Pistol" star, Sickle region	17:46:15.3	-28:50:04	3.2x3.2	6.3	0	0.047
"Pistol" star, Sickle region	17:46:15.3	-28:50:04	3.2x3.2	6.6	0	0.047
"Pistol" star, Sickle region	17:46:15.3	-28:50:04	3.2x3.2	7.7	0	0.042
"Pistol" star, Sickle region	17:46:15.3	-28:50:04	3.2x3.2	19.7	31.4	0.0083
"Pistol" star, Sickle region	17:46:15.3	-28:50:04	3.2x3.2	24.2	37.1	0.03
NGC 253	00:47:33.13	-25:17:17.8	3.2x3.2	5.4	0	0.5
NGC 253	00:47:33.13	-25:17:17.8	3.2x3.2	6.3	0	0.5
NGC 253	00:47:33.13	-25:17:17.8	3.2x3.2	6.6	0	0.5
NGC 253	00:47:33.13	-25:17:17.8	3.2x3.2	7.7	0	0.5
NGC 253	00:47:33.13	-25:17:17.8	3.2x3.2	8.6	0	0.5
NGC 253	00:47:33.13	-25:17:17.8	3.2x3.2	11.1	0	0.5
NGC 253	00:47:33.13	-25:17:17.8	3.2x3.2	11.3	0	0.5
NGC 253	00:47:33.13	-25:17:17.8	3.2x3.2	19.7	31.4	0.5
NGC 253	00:47:33.13	-25:17:17.8	3.2x3.2	24.2	37.1	0.5
M82	09:55:52.2	+69:40:47	3.2x3.2	5.4	0	0.41
M82	09:55:52.2	+69:40:47	3.2x3.2	8.6	0	0.037
M82	09:55:52.2	+69:40:47	3.2x3.2	11.1	0	0.046







M82	09:55:52.2	+69:40:47	3.2x3.2	19.7	0	0.0028
M82	09:55:52.2	+69:40:47	3.2x3.2	24.2	0	0.0076
NGC 4038/9	12:01:52	-18:52:02.9	3.2x3.2	19.7	31.4	1.13
NGC 4038/9	12:01:52	-18:52:02.9	3.2x3.2	19.7	37.1	1.13
NGC1068	02:42:40.83	-00:00:48.4	3.2x3.2	19.7	31.4	0.33
NGC1068	02:42:40.83	-00:00:48.4	3.2x3.2	19.7	37.1	0.33
Arp 299	11:28:32.8	+58:34:45	3.2x3.2	5.4	0	0.9
Arp 299	11:28:32.8	+58:34:45	3.2x3.2	6.3	0	3.64
Arp 299	11:28:32.8	+58:34:45	3.2x3.2	6.6	0	0.0014
Arp 299	11:28:32.8	+58:34:45	3.2x3.2	7.7	0	3.29
Arp 299	11:28:32.8	+58:34:45	3.2x3.2	11.1	0	2.08
Arp 299	11:28:32.8	+58:34:45	3.2x3.2	19.7	31.4	0.17
Arp 299	11:28:32.8	+58:34:45	3.2x3.2	19.7	37.1	0.17
M16	18:18:51.5	13:49:30	3.2x3.2	6.3	0	0.53
M16	18:18:51.5	13:49:30	3.2x3.2	6.6	0	2.16
M16	18:18:51.5	13:49:30	3.2x3.2	7.7	0	0.15
M16	18:18:51.5	13:49:30	3.2x3.2	0	33.5	0.91
M16	18:18:51.5	13:49:30	3.2x3.2	19.7	34.8	0.47
M16	18:18:51.5	13:49:30	3.2x3.2	24.2	37.1	1.17
M17	18:20:30	-16:10:11	9.6x9.6	6.3	0	1.18
M17	18:20:30	-16:10:11	9.6x9.6	6.6	0	1.18
M17	18:20:30	-16:10:11	9.6x9.6	7.7	0	1.07
M17	18:20:30	-16:10:11	9.6x9.6	0	33.5	0.15
M17	18:20:30	-16:10:11	9.6x9.6	19.7	34.8	0.07
M17	18:20:30	-16:10:11	9.6x9.6	24.2	37.1	0.16
W51	19:23:40	+14:31:08	6.4x6.4	6.3	0	0.26
W51	19:23:40	+14:31:08	6.4x6.4	6.6	0	0.26
W51	19:23:40	+14:31:08	6.4x6.4	7.7	0	0.23
W51	19:23:40	+14:31:08	6.4x6.4	0	33.5	0.093
W51	19:23:40	+14:31:08	6.4x6.4	19.7	34.8	0.048
W51	19:23:40	+14:31:08	6.4x6.4	24.2	37.1	0.12
Orion BNKL, Trapezium, Bar	05:35:16.0	-05::23:17	12.8x9.6	6.3	0	0.4
Orion BNKL, Trapezium, Bar	05:35:16.0	-05::23:17	12.8x9.6	6.6	0	0.4
Orion BNKL, Trapezium, Bar	05:35:16.0	-05::23:17	12.8x9.6	7.7	0	0.4
Orion BNKL, Trapezium, Bar	05:35:16.0	-05::23:17	12.8x9.6	0	31.4	0.4
Orion BNKL, Trapezium, Bar	05:35:16.0	-05::23:17	12.8x9.6	0	33.5	0.4
Orion BNKL, Trapezium, Bar	05:35:16.0	-05::23:17	12.8x9.6	19.7	34.8	0.4
Orion BNKL, Trapezium, Bar	05:35:16.0	-05::23:17	12.8x9.6	24.2	37.1	0.4
HV Tau	04:38:35.311	+26:10:38.49	3.2x3.2	19.7	31.4	0.43
HV Tau	04:38:35.311	+26:10:38.49	3.2x3.2	24.2	37.1	1.21
DK Tau	04:30:44.243	+26:01:24.79	3.2x3.2	19.7	31.4	0.043
DK Tau	04:30:44.243	+26:01:24.79	3.2x3.2	24.2	37.1	0.12
IT Tau	04:33:54.708	+26:13:27.70	3.2x3.2	19.7	31.4	0.43
IT Tau	04:33:54.708	+26:13:27.70	3.2x3.2	24.2	37.1	1.21







HK Tau	04:31:50.576	+24:24:17.84	3.2x3.2	19.7	31.4	0.11
HK Tau	04:31:50.576	+24:24:17.84	3.2x3.2	24.2	37.1	0.31
GG Tau B	04:32:30.326	+17 31 40.67	3.2x3.2	19.7	31.4	0.043
GG Tau B	04:32:30.326	+17 31 40.67	3.2x3.2	24.2	37.1	0.12
FX Tau	04:30:29.617	+24:26:45.04	3.2x3.2	19.7	31.4	0.17
FX Tau	04:30:29.617	+24:26:45.04	3.2x3.2	24.2	37.1	0.48
FQ Tau	04:19:12.798	+28:29:33.01	3.2x3.2	19.7	31.4	0.43
FQ Tau	04:19:12.798	+28:29:33.01	3.2x3.2	24.2	37.1	1.21
FV Tau	04:26:53.502	+26:06:54.06	3.2x3.2	19.7	31.4	0.043
FV Tau	04:26:53.502	+26:06:54.06	3.2x3.2	24.2	37.1	0.12
Haro 6-28	04:35:56.837	+22:54:36.22	3.2x3.2	19.7	31.4	0.86
Haro 6-28	04:35:56.837	+22:54:36.22	3.2x3.2	24.2	31.4	0.86
UZ Tau	04:32:42.962	+25:52:31.09	3.2x3.2	19.7	31.4	0.043
UZ Tau	04:32:42.962	+25:52:31.09	3.2x3.2	24.2	37.1	0.12
GH Tau	04:33:06.216	+24:09:33.72	3.2x3.2	19.7	31.4	0.17
GH Tau	04:33:06.216	+24:09:33.72	3.2x3.2	24.2	37.1	0.48
NGC 2024	05:41:43	-01:50:30	6.4x6.4	19.7	31.4	0.10
NGC 2024	05:41:43	-01:50:30	6.4x6.4	24.2	37.1	0.44
S106	20:27:27	37:22:48	6.4x3.2	19.7	31.4	0.66
S106	20:27:27	37:22:48	6.4x3.2	24.2	31.4	0.66
S106	20:27:27	37:22:48	6.4x3.2	0	37.1	1.45
Serpens	18:30:00.0	01:15:00	12.8x12.8	19.7	31.4	0.10
Serpens	18:30:00.0	01:15:00	12.8x12.8	0	37.1	0.089
L988-e	21:03:58	50:14:38	12.8x12.8	19.7	31.4	0.10
L988-e	21:03:58	50:14:38	12.8x12.8	0	37.1	0.089








Appendix A4 - FORCAST grism Reserved Observations Catalog (ROC)

Target	RA (J2000)	Dec (J2000)	S/N	Obs time/ slit pos [h]	Obs time [h]	Map Area	Mode	Grism [µm]
Orion A, BN/KL	05:35:14.16	-05:22:21.5	20	0.25	1	3'x3'	Long slit	5-8, 8-13.7, 17-28, 28-40
Orion A, BN/KL	05:35:14.16	-05:22:21.5	20	0.5	2	30"x30"	Short slit /high res.	5-8, 8-13.7
Orion A Bar North	05:32:56.0	-05 26:30.0	20	3.0	6.0	3.2'x3.2'	Long slit	5-8, 8-13.7, 17-28, 28-40
Orion A, Bar South	05:32:49.0	-05:27:15.0	20	3.0	6.0	3.2'x3.2'	Long slit	5-8, 8-13.7, 17-28, 28-40
Sgr A*	17:42:29.31	-28:59:18.5	40	0.125	1.25	3.2'x3.2'	Long slit	5-8, 8-13.7, 17-28, 28-40
AFGL 2688	21:00:19.37	+36:29:44.3	20	0.125	0.25	±10" EW	Long slit	5-8, 8-13.7, 17-28, 28-40
AFGL 2688	21:00:19.37	+36:29:44.3	10	0.125	0.25	±10" EW	Short slit /high res.	5-8, 8-13.7








Appendix A5 - GREAT Cycle 3 Reserved Observations Catalog (ROC)

Science	Object Name	RA	DEC	ν1	ν 2	V 3	area	Time
		(2000)	(2000)		[THz]		arcmin	[hr]
PP disks	DG Tau	04:27:04.7	+26:06:16.3	all	all lines below		0.3	0.5
	HL Tau	04:31:38.4	+18:13:57.7	all lines below		W	0.3	0.5
	AB Aur	04:55:45.8	-06:57:59.5	all lines below		0.3	0.5	
	HD50138	06:51:33.4	-06:57:59.5	all lines below		0.3	0.5	
	HD100546	11:33:25.4	-70:11:41.2	all	lines belo)W	0.3	0.5
	HD163296	17:56:21.3	-21:57:21.9	all	lines belo)W	0.3	0.5
	HD97048	11:08:03.3	-77:39:17.4	all	lines belo)W	0.3	0.5
Star formation	IRAS16293-2422	16:32:22.6	-24:28:33.0	#1	#2	4.7	1	done
cores	SgrB2(M)(N)	17:47:20.4	-28:23:07.0	#1	#2	4.7	2	done
	W28A	18:00:30.4	-24:04:00.0	#1	#2	4.7	1	0.3
	W31C	18:10:28.7	-19:55:50.0	#1	#2	4.7	1	0.3
	W33A	18:14:39.4	-17:52:00.0	#1	#2	4.7	1	0.3
	W49N	19:10:13.2	+09:06:12.0	#1	#2	4.7	1	done
	DR21(OH)	20:39:00.7	+42:22:46.7	#1	#2	4.7	2	0.6
	W43-MM1	18:47:47.0	-01:54:28.0	#1	#2	4.7	1	0.3
	W3- IRS1	02:25:40.6	+62:05:51.0	#1	#2	4.7	1	0.6
	W3(OH)	02:27:04.1	+61:52:22.0	#1	#2	4.7	1	0.6
	G31.41	18:47:34.3	-01:12:46.0	#1	#2	4.7	1	0.3
	G29.96	18:46:03.8	-02:39:22.0	#1	#2	4.7	1	0.3
	NGC7538	23:13:45.3	+61:28:10.0	#1	#2	4.7	2	0.6
	W51D/E1	19:23:43.8	+14:30:26.0	#1	#2	4.7	2	0.6
	G34.26	18:53:18.6	+01:14:58.0	#1	#2	4.7	1	done
	G10.47	18:08:38.2	-19:51:50.0	#1	#2	4.7	1	0.3
	G5.89	18:00:30.4	-24:04:02.0	#1	#2	4.7	1	0.3
	Orion-KL	05:35:15.1	-05:22:26.6	#1	#2	4.7	2	0.5
	Orion-Bar	05:35:22.1	-05:25:13.4	#1	#2	4.7	2	0.6
	M17SW	18:20:27.6	-16:12:00.9	CO	#2	4.7	1	done
	G333.6	16:22:09.0	-50:06:15.0	CO	#2	4.7	30	1.0
	NGC2024	05:41:45.2	-01:55:45.0	CO	#2	4.7	2	1.0
	S106	20:27:26.7	37:22:47.9	CO	#2	4.7	2	1.0
	NGC6334I	17:20:53.3	-35:47:01.5	#1	#2	4.7	2	0.5
	NGC6334I(N)	17:20:55.1	-35:45:08.8	#1	#2	4.7	2	0.5
	IRAS17233	17:26:42.5	-36:09:18.1	#1	#2	4.7	2	1.0
	AG351.58-0.35	17:25:25.0	-36:12:45.4	#1	#2	4.7	2	0.5







p-H ₂ D ⁺ /o-D ₂ H ⁺	IRAS16293E	16:32:29.4	-24:28:52.6	1.37		1.48	1	1
p-1120 10-0211	IRAS16293A/B	16:32:22.9	-24:28:39.7	1.37		1.48	1	0.5
0 (0) - ()	IRAS18151-1208	18:18:10.3	-12:07:27.0	CO	#2	4.7	1	0.5
Outflow studies	IRAS17233-3606	17:26:42.5	-36:09:18.0	CO	#2	4.7	1	0.5
	IRAS05358+3543	05:39:13.1	+35:45:50.0	CO	#2	4.7	1	0.5
	IRAS20126+4104	20:14:25.1	+41:13:32.0	CO	#2	4.7	1	0.5
	eta Carina	10:45:03.6	-59:41:04.3	CO	#2	4.7	1	0.5
	HH212	05:43:51.4	-01:02:53.0	CO	#2	4.7	2	0.7
	BHR71	12:01:36.3	-65:08:53.0	CO	#2	4.7	2	done
	HH111	05:51:46.3	+02:48:30.0	CO	#2	4.7	2	0.7
	NGC2071	05:47:04.7	+00:21:44.0	CO	#2	4.7	2	1.0
	NGC1333	03:28:55.6	+31:14:37.1	CO	#2	4.7	3	1.0
	HH54	12:55:50.3	-76:56:23.0	CO	#2	4.7	2	0.5
	G327-0.6	15:53:08.8	-54:37:01.0	CO	#2	4.7	2	0.5
	AG012.81-0.20	18:14:13.5	-17:55:32.0	CO	#2	4.7	2	0.5
	AG015.03-0.67	18:20:22.4	-16:11:43.8	CO	#2	4.7	3	0.5
Stars	ALF-Ori	05:55:10.3	+07:24:25.0	СО	1.9	4.7	1	0.5
	VY-CMa	07:22:58.3	-25:46:03.1	CO	1.9	4.7	1	0.5
	O-Ceti	02:19:20.8	-02:58:43.0	CO	1.9	4.7	1	0.5
	IK-Tau	03:53:28.9	+11:24:21.7	CO	1.9	4.7	1	0.5
(P)PNe	NGC7027	21:07:01.6	+42:14:10.2	CO	1.9	4.7	1	done
	CRL618	04:42:53.6	+36:06:53	CO	1.9	4.7	0.5	0.5
	CRL2688	21:02:18.7	+36:41:37.8	CO	1.9	4.7	0.5	0.5
	OH231.8+4.2	07:42:16.8	-14:42:52.1	CO	1.9	4.7	0.5	0.5
	NGC6302	17:13:29.3	-37:06:11.2	CO	1.9	4.7	0.5	0.5
	BD+30 3639	19:34:45.2	+30:30:58.8	CO	1.9	4.7	0.5	0.5
	IRAS21282+5050	21:59:58.4	+51:03:59.8	CO	1.9	4.7	0.5	0.5
	IC1396	21:40:42.3	+58:16:10.0	СО	1.9	4.7	60	1.0
Galactic PDRs	DR21C	20:39:00	+42:19:42.0	CO	1.9	4.7	4	done
	IC63	00:58:55	+60:53:11.0	CO	1.9	4.7	2	1.0
	Carina N	10:43:30.9	-59:34:16.3	CO	1.9	4.7	3	2.0
	NGC3603	11:15:08.9	-61:16:50.0	СО	1.9	4.7	5	1.0
	OMC2	05:35:25.0	-05:12:30.0	CO	1.9	4.7	7	1.5
	OMC3	05:12:30.0	-05:03:45.0	CO	1.9	4.7	7	1.5
	Orion-fSE1/2	05:36:15.0	-05:38:00.0	CO	1.9	4.7	15	2.0







LMC PDRs	N159	05:39:57.2	-69:44:33.0	1.46	1.9	4.7	2	1.0
	30 Dor	05:38:42.4	-69:06:03.3	1.46	1.9	4.7	2	1.0
	N160	05:39:40.3	-69:39:01.1	1.46	1.9	4.7	1	1.0
SMC PDRs	N88	01:24:08.5	-73:08:53.2	1.46	1.9	4.7	1	0.5
	N66	00:59:27.4	-72:10:10.7	1.46	1.9	4.7	1	0.5
	Sickle	17:46:14.7	-28:48:55.9	СО	1.9	4.7	3	0.75
Galactic Center	Arches	17:45:51.0	-28:48:22.0	CO	1.9	4.7	3	0.75
Studies	M-002-007	17:45:51.7	-28:59:13.7	CO	1.9	4.7	3	0.5
	M-013-008	17:45:36.8	-29:05:23.8	CO	1.9	4.7	3	0.5
	SgrA /CND	17:45:39.9	-29:00:28.2		all bands		10	2.0
	M+0.253	17:46:09.0	-28:42:40.0	CO	1.9	4.7	3	1.0
	M-3.8+0.9	17:32:53.5	-31:43:21.8	CO	1.9	4.7	10	1.0
Chlorine	Venus	HCI		1.25		4.7	_	0.75
chemistry	Jupiter	HCI		1.25		4.7	_	0.75
,	Saturn	HCI		1.25		4.7	-	0.75
Hydrogen		HDO, (H ₂ ¹⁶ O), H ₂ ¹⁸ O		1.27	1.89	4.7	-	0.75
chemistry	Mars	H ₂ O ₂			1.85	1.90	-	0.75
		1					1	
nearby nuclei	M82	09:55:52.2	+69:40:49.6		lines belo		1	2.0
	NGC253	00:47:33.1	-25:17:17.6	all	lines belo)W	1	1.5
	NGC4945	13:05:27.5	5 -49:28:05.6		all lines below		1	1.5
	Cen-A	13:25:27.6	-43:01:08.9			4.7	1	1.5
MHD shocks	IC443	06:17:42.5	+22:21:30.0	all lines below		40	2.0	
MILE SHOCKS	W28F	18:01:52.3	-23:19:25.0	CO	#2	4.7	2	done
	W44 E/F	18:56:28.4	+01:29:55.0	CO	#2	4.7	2	1.0
	3C391	18:49:22.3	-00:57:22.0	CO	#2	4.7		1.0

Times given in the last column are total integration times (on and off source), but no overheads due to calibration/facility inefficiencies have been added. Min map size of observation is one arcmin (though in most cases this will be the central beam only). In most targets "CO" refers to the J-transition accessible in bands L1-L2, including selected isotopologues.

Frequency (THz) Species

set-up #2 (in band L2): 1.81 (NH3), #2 (OH), 1.9 ([CII])

set-up #1 (in band L1): 1.39 (OD), CO


1.37 $p-H_2D^+$

1.39 OD

1.46 [N II]

1.48 $\text{o-D}_2\text{H}^+$

4.7 [OI]







Appendix A6 - FIFI-LS Reserved Observation Catalog

Target	RA (J2000)	DEC (J2000)	Extent (arcmin)	Lines B λ in μm	Lines R λ in μm	Time (h)
M42	05:35:16.6	-05:23:28.1	3 x 5	[OIII] \(\lambda 52\) [OIII] \(\lambda 88\)	CO (17-16) λ153 OH λ163	3
M42 completing existing map	05:35:16.6	-05:23:28.1	6 x 7	[OI] \(\lambda 63\) CO(22-21) \(\lambda 119\)	[CI] λ157 [OI] λ145	1
Horsehead	05:40:59.0	-02:27:30	2 x 4	[OIII] \\ \text{\lambda}52 \\ [OII] \\ \text{\lambda}88 \\ [OI] \\ \text{\lambda}63 \end{array}	[CII] λ157 [OI] λ145 CO (17-16) λ153	3
Brick	17:46:08.6	-28:42:46.0	3 x 5	[OI] λ63 OH λ79	[OI] λ145 [CII] λ157	3
M17-SW	18:20:23.1	-16:11:43	2.5 x 4	[OIII] \\ \text{\lambda}52 \\ [OIII] \\ \text{\lambda}88 \\ [OI] \\ \text{\lambda}63 \end{array}	[CII] λ157 [OI] λ145 CO (17-16) λ153	3
W43-main	18:47:40.0	-01:57:00.0	5 x 5	[OIII] \\(\lambda 52\) [OIII] \\(\lambda 88\) OH \(\lambda 79\)	[OI] λ145 [CII] λ157 CO (14-13) λ186	3
DR21(OH)	20:39:00.7	+42:22:46.7	1 x 1	[OIII] \\ \text{\lambda}52 \\ [OIII] \\ \text{\lambda}88 \\ OH \\ \text{\lambda}79 \end{array}	[OI] λ145 [CII] λ157	1
SgrA*	17:45:40	-29:00:28	2.5 x 2.5	OH λ79 OH λ119 [NII] λ57	CO (15-14) λ174 CO (19-18) λ137 CO (20-19) λ130	2
SgrA*	17:45:40	-29:00:28	10 x 10	[OI] λ63 [OIII] 88 OH 119	[CII] λ157 [OI] λ145 CO (14-13) λ186	6
Sickle	17:46:12	-28:48:30	3 x 3	[OIII] \(\lambda 52\) [OII] \(\lambda 88\) [OI] \(\lambda 63\)	[CII] λ157 [OI] λ145 CO (17-16) λ153	2
Arches	17:45:47	-28:50:40	1 x 1.5	[OIII] \\ \text{\lambda}52 \\ [OIII] \\ \text{\lambda}88 \\ [OI] \\ \text{\lambda}63 \end{array}	[CII] λ157 [OI] λ145 CO (17-16) λ153	1
NGC 253	00:47:33.1	-25:17:18	2 x 2	[OIII] \\(\lambda 52\) [OIII] \\(\lambda 88\) [OI] \\(\lambda 63\)	[CII] λ157 [OI] λ145 CO(14-13) λ186	1.5
M82	09:55:52.4	+69:40:47	2 x 2	[ΟΙΙΙ] λ52	[CII] λ157	2
M83	13 37 00.9	-29 51 57	3 x 3	[OIII] \\(\lambda 52\) [OII] \\(\lambda 88\) [OI] \\(\lambda 63\)	[CII] λ157 [OI] λ145 CO(14-13) λ186	5
IC10	00:20:17.3	+59:18:13.6	5 x 5	[OIII] λ52 [NIII] λ57	[CII] λ157 [OI] λ145	5
NGC1140	02:54:33.6	10:01:39.9	2x2	[OIII] 52	[OI] λ145	4
He2-10	8:36:15.18	-26:24:33.9	1 x 1	[OIII] 52	[OI] λ145 CO (14-13) λ186	3
NGC4449	12:28:11.12	+44:05:36.8	2 x 2	[OIII] λ52 [NIII] λ57	[ΟΙ] λ145	4
NGC5253	13:39:55.96	-31:38:24.4	1x1	[OIII] λ52 [NIII] λ57	[OI] λ145	4

