

## THE SPITZER-HETDEX EXPLORATORY LARGE-AREA SURVEY

C. PAPOVICH<sup>1,2</sup>, H. V. SHIPLEY<sup>1,2</sup>, N. MEHRTENS<sup>1,2</sup>, C. LANHAM<sup>1</sup>, M. LACY<sup>3</sup>, S. L. FINKELSTEIN<sup>4</sup>, R. BASSETT<sup>5</sup>, P. BEHROOZI<sup>6</sup>, G. BLANC<sup>7,8,9</sup>, R. CIARDULLO<sup>10,11</sup>, R. DE JONG<sup>12</sup>, D. DEPOY<sup>1,2</sup>, N. DRORY<sup>4</sup>, E. GAWISER<sup>13</sup>, K. GEBHARDT<sup>4</sup>, C. GRONWALL<sup>10,11</sup>, G. HILL<sup>4</sup>, U. HOPP<sup>14</sup>, S. JOGEE<sup>4</sup>, L. KAWINWANICHAKIJ<sup>1,2</sup>, J. MARSHALL<sup>1,2</sup>, E. MCLINDEN<sup>4</sup>, E. MENTUCH COOPER<sup>4</sup>, R. SOMERVILLE<sup>13</sup>, M. STEINMETZ<sup>12</sup>, K.-V. TRAN<sup>1,2</sup>, S. TUTTLE<sup>4</sup>, M. VIERO<sup>15</sup>, R. WECHSLER<sup>15</sup>, G. ZEIMANN<sup>4,10,11</sup>,

<sup>1</sup>Department of Physics and Astronomy, Texas A&M University, College Station, TX, 77843-4242 USA; papovich@physics.tamu.edu

<sup>2</sup>George P. and Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, College Station, TX, 77843-4242 USA

<sup>3</sup>North American ALMA Science Center, NRAO Headquarters, Charlottesville, VA 22903, USA

<sup>4</sup>Department of Astronomy, The University of Texas at Austin, Austin, TX 78712, USA

<sup>5</sup>Centre for Astrophysics & Supercomputing, Swinburne University, Hawthorn, VIC 3122, Australia

<sup>6</sup>Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218

<sup>7</sup>Departamento de Astronomía, Universidad de Chile, Camino del Observatorio 1515, Las Condes, Santiago, Chile

<sup>8</sup>Centro de Astrofísica y Tecnologías Afines (CATA), Camino del Observatorio 1515, Las Condes, Santiago, Chile

<sup>9</sup>Visiting Astronomer, Observatories of the Carnegie Institution for Science, 813 Santa Barbara St, Pasadena, CA 91101, USA

<sup>10</sup>Department of Astronomy and Astrophysics, The Pennsylvania State University, University Park, PA 16802, USA

<sup>11</sup>Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, PA 16802, USA

<sup>12</sup>Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, D-14482 Potsdam, Germany

<sup>13</sup>Department of Physics and Astronomy, Rutgers, The State University of New Jersey, 136 Frelinghuysen Road, Piscataway, NJ 08854, USA

<sup>14</sup>Max-Planck-Institut für Extraterrestrische Physik, D 85741, Garching, Germany and

<sup>15</sup>Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, 382 Via Pueblo Mall, Stanford, CA 94305, USA

To be submitted to *ApJS*, Draft Version October 14, 2015

### ABSTRACT

We present post-cryogenic *Spitzer* imaging at 3.6 and 4.5  $\mu\text{m}$  with the Infrared Array Camera (IRAC) of the *Spitzer*/HETDEX Exploratory Large-Area (SHELA) survey. SHELA covers  $\approx 24 \text{ deg}^2$  of the Sloan Digital Sky Survey “Stripe 82” region, and falls within the footprints of the Hobby-Eberly Telescope Dark Energy Experiment (HETDEX) and the Dark Energy Survey. The HETDEX blind  $R \sim 800$  spectroscopy will produce  $\sim 200,000$  redshifts from the Lyman- $\alpha$  emission for galaxies in the range  $1.9 < z < 3.5$ , and an additional  $\sim 200,000$  redshifts from the [O II] emission for galaxies at  $z < 0.5$ . When combined with deep *ugriz* images from the Dark Energy Camera, *K*-band images from NEWFIRM, and other ancillary data, the IRAC photometry from *Spitzer* will enable a broad range of scientific studies of the relationship between structure formation, galaxy stellar mass, halo mass, AGN, and environment over a co-moving volume of  $\sim 0.5 \text{ Gpc}^3$  at  $1.9 < z < 3.5$ . Here, we discuss the properties of the SHELA IRAC dataset, including the data acquisition, reduction, validation, and source catalogs. Our tests show the images and catalogs are 80% (50%) complete to limiting magnitudes of 22.0 (22.6) AB mag in the detection image, which is constructed from the weighted sum of the IRAC 3.6 and 4.5  $\mu\text{m}$  images. The catalogs reach limiting sensitivities of 1.1  $\mu\text{Jy}$  at both 3.6 and 4.5  $\mu\text{m}$  ( $1\sigma$ , for  $R = 2''$  circular apertures). As a demonstration of science, we present IRAC number counts, examples of highly temporally variable sources, and galaxy surface density profiles of rich galaxy clusters. In the spirit of *Spitzer* Exploratory programs we provide all images and catalogs as part of the publication.

**Keywords:** catalogs — galaxies: clusters: general — infrared: galaxies — surveys


### 1. INTRODUCTION

The launch of the *Spitzer* Space Telescope (Werner et al. 2004) allowed for large surveys of galaxies at near-IR wavelengths, which are free from foreground terrestrial thermal emission and are sensitive to the rest-frame peak of the stellar emission in galaxies ( $\lambda_{\text{rest}} \sim 1.6 \mu\text{m}$ ) over redshifts  $z \sim 1-2$  (e.g., Eisenhardt et al. 2008; Papovich 2008; Muzzin et al. 2013). During the cryogenic mission, *Spitzer* executed a variety of initial, wide-area surveys (e.g., Lonsdale et al. 2003; Ashby et al. 2009; Wilson et al. 2009), and the post-cryogenic (“warm”) mission enabled much larger surveys with increasingly larger combinations of depth and area (Mauduit et al. 2012; Ashby et al. 2013a,b, 2015; Labbé et al. 2013; Timlin et al. 2015).

The size and depth of the near-IR imaging surveys carried out by *Spitzer* have expanded our knowledge of how dark matter halos accumulate baryons and convert them into stars. The physics governing this formation involves a range of com-

plex processes (see the recent review by Somerville & Davé 2015, and references therein). The processes for the growth of galaxies include baryon and dark-matter accretion histories, gas cooling, star formation, and galaxy mergers, while the processes that inhibit this growth include energetic winds from massive stars and supernovae, radiation and kinematic feedback from AGN, and shock heating of gas in large dark matter halos. Distinguishing the importance of these different processes, and their dependence on halo and stellar mass, redshift, and environment is one of the main goals of galaxy formation theory.

*Spitzer* has allowed us to test some of the theoretical physical processes by comparing measurements of the galaxy stellar mass distribution with model predictions. All of these theoretical physical processes are expected to manifest as a function of galaxy stellar mass, halo mass, redshift, and environment. By connecting galaxies to their halos, we can identify and constrain the relative importance of the physical processes responsible for galaxy growth at different stages of



**Figure 1.** Field layout for SHELA. The thick lines show the coverage of our SHELA IRAC data (thick red-lined region), the SDSS Stripe 82 coverage (white-lined region), the planned coverage of the full-field of the HETDEX survey (yellow-lined region), the HerS *Herschel* coverage (magenta-lined region) and the planned coverage of the DES (green-lined region). The lines of constant R.A. and Decl. are labeled. The background image shows the IRAS 100  $\mu\text{m}$  map of a portion of the south Galactic Pole (Schlegel et al. 1998). The image intensity scales with surface brightness as indicated in the color bar in units of MJy/sr.

their evolution. Recent studies have attempted this by measuring the ratio of galaxy stellar mass to halo mass (SM–HM) as a function of halo mass (e.g., Moster et al. 2010, 2013; Behroozi et al. 2013; Kravtsov et al. 2014). The SM–HM relation provides a powerful mechanism for connecting the predictions for the halo mass function (which is well understood, e.g., Springel et al. 2005; Tinker et al. 2008; Behroozi et al. 2010, 2013) to the observed stellar mass functions and mass-dependent spatial clustering of galaxies (e.g., Weinberg et al. 2004).

Expanding tests of galaxy formation derived from the SM–HM relation requires observational measurements of galaxies over large areas to measure the both the bulk statistics and scatter in the halo- and stellar-mass distributions. This drives the need for larger extragalactic surveys that cover large areas containing the full range of environments in which galaxies form, and during the epochs when the physical processes manifest.

The post-cryogenic *Spitzer* mission has enabled such surveys. Here, we describe one such program, the *Spitzer*/HETDEX Exploratory Large Area (SHELA) survey, which is designed to measure the evolution of the nature of the SM–HM relation for galaxies over a large baseline in redshift,  $1.9 < z < 3.5$ . SHELA targets a  $\approx 24 \text{ deg}^2$  field in the Sloan Digital Sky Survey (SDSS) Stripe 82 field (Annis et al. 2014), and covers a portion of the footprint of the Hobby-Eberly Telescope (HET) Dark Energy eXperiment (HETDEX, Hill et al. 2008). The SHELA field contains a large amount of ground-based optical and near-IR imaging, including forthcoming optical imaging data from the Dark Energy Survey (DES), as well as coverage at far-IR, sub-mm, and X-ray wavelengths. The large SHELA field covers nearly  $0.5 \text{ Gpc}^3$  in cosmological volume at both moderate redshifts,  $0.5 < z < 2.0$ , and at high redshifts,  $2.0 < z < 3.5$ , and opens the distant Universe in the way that large-area, shallow surveys have for the low-redshift Universe (such as the Sloan Digital Sky Survey, SDSS, see Alam et al. 2015). As an equatorial field, the SHELA field is accessible to terrestrial tele-

scopes in both hemispheres, which gives it a high and lasting legacy value for studies of galaxy evolution, AGN, and large-scale structure.

Here we present the overview of the *Spitzer* imaging dataset for SHELA. The outline for this *Paper* is as follows. In § 2, we describe the SHELA survey field and the survey strategy with the *Spitzer* space telescope. In § 3, we describe the data reduction and mosaicking of the *Spitzer* dataset, and we describe astrometric and photometric quality checks on the imaging data. In § 4, we discuss the construction of the source catalog, and the catalog properties, including catalog completeness. We also discuss estimates of photometric errors. In § 5, we discuss basic scientific results, including source number counts, temporally varying objects, and the galaxy surface density of rich clusters. In § 6, we summarize the work. Throughout, we denote photometric magnitudes measured in the IRAC channel 1 and channel 2 as [3.6] and [4.5], respectively. Unless stated otherwise, all magnitudes here are relative to the AB system (Oke & Gunn 1983), where  $m_{\text{AB}} = 23.9 - 2.5 \log(f_{\nu}/\mu\text{Jy})$ . For convenience, we provide conversions between the AB system and the system relative to Vega,  $[3.6]_{\text{AB}} - [3.6]_{\text{Vega}} = 2.79 \text{ mag}$  and  $[4.5]_{\text{AB}} - [4.5]_{\text{Vega}} = 3.26 \text{ mag}$ , derived from a comparison to the spectrum of an A0V spectral type star. Users of the catalog may apply these to the flux densities in the catalog to convert them to the magnitude system relative to Vega. For any derived, physical quantity, we assume a cosmology with  $\Omega_m = 0.3$ ,  $\Omega_{\Lambda} = 0.7$ , and  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ , consistent with the WMAP seven-year data (Komatsu et al. 2011) and Planck 2013 data (Planck Collaboration et al. 2014).

## 2. FIELD AND SURVEY CHARACTERIZATION

### 2.1. HETDEX

SHELA covers a portion of the equatorial field of HETDEX. HETDEX is a dark energy study which will produce the most significant constraint on the evolution of dark energy from  $z \sim 2$  to the present. HETDEX is a spectroscopic survey, which will measure redshifts of Lyman- $\alpha$  emitting galax-

**Table 1**  
Observation Log for IRAC Observations

Observing Epoch (1)	Observing Dates (UTC) (2)	Position Angles (deg. E of N) (3)	Number of AORS (4)
1	2011-09-28 to 2011-10-10	-110.0 to -105.5	64
2	2012-02-13 to 2012-02-29	63.1 to 65.3	63
3	2012-09-28 to 2012-10-09	-110.9 to -107.4	64

ies (LAEs) between  $1.9 < z < 3.5$  using a suite of up to 78 wide-field integral field units (IFUs) covering 350–550 nm, simultaneously (Hill et al. 2008). The entire HETDEX survey will cover 420 deg<sup>2</sup> with a 1/4.5 filling factor with blind pointings, detecting  $0.8 \times 10^6$  LAEs and more than  $10^6$  [O II]-emitting galaxies with  $z < 0.5$ . HETDEX has two planned fields: a high Galactic declination field with 60 deg<sup>2</sup> of sky with spectra, and an equatorial field with 30 deg<sup>2</sup> with spectra.<sup>2</sup> For the SHELA field, our plans are to increase the filling factor to unity such that the entire field has spectroscopic coverage to the HETDEX limit ( $10\sigma$  limits for line flux of  $3.4 \times 10^{-17}$  erg cm<sup>-2</sup> s<sup>-1</sup> or continuum of  $g_{AB}=21.9$  mag).


SHELA covers a  $\approx 24$  deg<sup>2</sup> field with IRAC coverage in the footprint of the HETDEX equatorial field. Figure 1 shows the location of the equatorial HETDEX field and the SHELA field. The SHELA/IRAC imaging will detect  $>200,000$  galaxies at  $1.9 < z < 3.5$  down to a limiting stellar mass of  $\approx 2-3 \times 10^{10} M_{\odot}$  (both LAEs and non-LAEs; assuming the  $3\sigma$  SHELA IRAC flux limit, for a stellar population formed at  $z_f = 5$ ). The goal for the combined HETDEX–SHELA dataset is to measure the relation between halo-mass (constrained by the HETDEX density field) and stellar-mass (constrained by the optical/IR colors). The size of the HETDEX/SHELA dataset ensures the analysis will be robust against statistical uncertainties.

Results from the HETDEX pilot survey using a HETDEX IFU on the McDonald 2.7m illustrate the impact of joint spectroscopy and stellar-population modeling of the LAE populations in small-area fields where optical/near-IR imaging and *Spitzer*/IRAC imaging already exist. These include results published in Adams et al. (2011), Blanc et al. (2011), Finkelstein et al. (2011), and Hagen et al. (2014). The results of the pilot survey also give us confidence that we understand the properties of our LAE selection, including their luminosity function and our ability to select LAEs for HETDEX with little contamination.

## 2.2. Field Location and Ancillary Data

The SHELA field covers a region within the HETDEX equatorial field. Figure 1 shows the coverage on the sky for these fields. The SHELA field is centered at R.A. =  $1^h 22^m 00^s$ , Decl. =  $+00^{\circ} 00' 00''$  (J2000), (Galactic coordinates,  $l = 138.294^{\circ}$ ,  $b = -62.017^{\circ}$ ) and extends approximately  $\pm 6.5$  deg in R.A. and  $\pm 1.25$  deg in Decl. The field was chosen to have low IR background (Schlegel et al. 1998) within the SDSS Stripe 82 and DES fields. As illustrated in Figure 1, the  $100 \mu\text{m}$  background ranges from 1.2 to 1.7 MJy/sr across the field, and with mean value of approximately 1.5 MJy/sr.

Because of its equatorial location, the SHELA field lies near the Ecliptic ( $\lambda = 18.93^{\circ}$ ,  $\beta = -8.01^{\circ}$ , and ranges in lat-



**Figure 2.** Example of persistence in the IRAC 3.6  $\mu\text{m}$  data. The data are from AOR 42817792 of SHELA. Each panel shows the same region of the IRAC detector in a time series of sequential BCDs, from 0061–0066. In the first three BCDs (0061–0063), the bright star (HD 9670,  $V = 6.9$  mag) falls at position 1, 2, and 3 in this portion of the IRAC detector. The bright star leaves a fractionally small persistence effect ( $< 0.01\%$  of the fluence) in the same x,y pixels in subsequent BCDs (0064–0066), which decays with an exponential timescale.

itude from  $\beta = -4^{\circ}$  to  $-11^{\circ}$ ). Because the primary component of the background for *Spitzer*/IRAC is the Zodiacal light, this results in a higher background than higher (Ecliptic) latitude fields. The Ecliptic latitude for SHELA falls between the values assumed for the “medium” and “high” background in the *Spitzer* sensitivity performance estimation tool (SENS-PET)<sup>3</sup>. Therefore, it is expected that the SHELA field will suffer higher-than average Zodiacal backgrounds, which adversely affects the flux sensitivity of the IRAC data.


An advantage of the equatorial location is that the SHELA field is readily observable by current and future optical/IR and radio telescopes. The SHELA field is centered on the equator, and overlaps with the DES optical imaging, and the optical imaging from the deeper SDSS/Stripe 82 coadd (Annis et al. 2014). These data are supplemented with our own deeper CTIO/DECam *ugriz* data, which reaches  $5\sigma$  limiting magnitudes of  $u = 25.8$ ,  $g = 25.8$ ,  $r = 25.3$ ,  $i = 25.0$ , and  $z = 24.5$  (in  $2''$ -diameter apertures). In addition, the field is being imaged in the  $K_s$  band down to a  $5\sigma$  depth of 22.8 mag using the NEWFIRM camera at Kitt Peak (PI: S. Finkelstein). The SHELA field has 250, 350, and 500  $\mu\text{m}$  images from the SPIRE instrument taken as part of the *Herschel* Stripe 82 Survey (HerS, Viero et al. 2014), and X-ray coverage from *Chandra* and *XMM-Newton* (LaMassa et al. 2013a,b). Finally, the SHELA field has received microwave observations at 148, 218, and 270 GHz from the Equatorial Survey of the Atacama Cosmology Telescope (ACT, Hasselfield et al. 2013). The growing amount of multiwavelength data make SHELA a rich survey for galaxy evolution and for studies of galaxy properties as a function of the large-scale structure.

Additional *Spitzer*/IRAC imaging of the SDSS Stripe 82 field exists from the *Spitzer* IRAC Equatorial Survey (SpIES, proposal ID (PID) 90045, PI: G. Richards; Timlin et al. 2015). The SpIES data cover an additional  $\sim 115$  deg<sup>2</sup> outside the SHELA footprint along SDSS Stripe 82, with an effective IRAC integration time of 120 s. Scaling by integration times, the SHELA data are approximately  $2.5 \log(\sqrt{270}/120) = 0.44$  mag deeper than SpIES. The reader is referred to Timlin et al., for a description of SpIES and its data products.

<sup>2</sup> The HETDEX equatorial field covers 150 deg<sup>2</sup> on the sky, but with a 1/4.5 fill factor, which makes the area covered with spectra closer to the 30 deg<sup>2</sup> quoted here.

<sup>3</sup> <http://ssc.spitzer.caltech.edu/warmmission/propkit/pet/senspet>





**Figure 3.** Combined, three-epoch mosaic of the SHELA IRAC Channel 1 ( $3.6 \mu\text{m}$ ) data. The extreme edges of the image cover nearly  $2.5 \text{ deg} \times 13 \text{ deg}$ , but the area covered to our 3-epoch depth is  $\approx 24 \text{ deg}^2$ .

### 2.3. SHELA IRAC Survey Strategy

In the post-cryogenic mission, IRAC (Fazio et al. 2004b) observes simultaneously in channel 1 (at  $3.6 \mu\text{m}$ ) and channel 2 (at  $4.5 \mu\text{m}$ ). Each channel observes a  $5'2 \times 5'2$  field. The field centers observed by each channel are separated on the sky by approximately 6.7 arcminutes, with a gap of about 1.52 arcminutes between the fields. We designed the mapping strategy for SHELA to obtain coverage in both channels over approximately the same area of sky.


Several constraints affected the design of our survey. We desired multiple dithers with slightly different position angles to allow redundancy, to identify cosmic rays, and to guard against image defects. We desired observations during several epochs to ease *Spitzer* scheduling requirements. The multiple epochs are separated by long enough periods of time (approximately 4–7 months) to identify time-variable objects including asteroids (as our observations are close to the ecliptic we expect asteroids to be detected at higher rates than higher-latitude extragalactic fields). We also required that all astronomical observation requests (AORs) be shorter than the maximum observing time, about 6 hours for *Spitzer*.

We divided the SHELA observations into three epochs, sep-

arated by approximately six months. There were two, 30-day duration observing windows each year for *Spitzer* to observe SHELA at position angles optimal for our survey strategy.

During each epoch we observed the entire SHELA field to one-third of the total depth, covering approximately  $12 \times 2.5 \text{ deg}^2$ . Each AOR used a three point dither pattern, with  $1 \times 30 \text{ s}$  frame time per position (where the array observes with 23.6 s of exposure time for a 30 s frame). Each AOR obtained a map divided into 8 rows by 10 columns of IRAC pointings, with a step size of  $280''$  between each pointing. The area covered by each AOR is approximately  $38' \times 47'$ , and each epoch tiled the entire SHELA field using 64 AORS (epochs 1 and 3) or 63 AORS (epoch 2). A single AOR required approximately 2.75 hrs of clock time. As there are 191 AORS, the total clock time for SHELA required 525 hrs of *Spitzer* observations. The *Spitzer* observations of SHELA occurred in the three epochs using these AORS under program PID 80100 (PI: Papovich), with dates listed in Table 1. The table also gives the position angles of IRAC during the observations and the number of AORS observed during each epoch.





**Figure 4.** Same as Figure 3, but for the SHELA IRAC Channel 2 ( $4.5 \mu\text{m}$ ) data.

### 3.1. Data Reduction

The SHELA IRAC data reduction began with the data pipeline processing version S19.0.0 of the IRAC Corrected Basic Calibrated Data (cBCDs) provided by the *Spitzer* Science Center (SSC). The data processing includes a subtraction of the dark current and application of the flat field, as well as a photometric and astrometric calibration. Starting with the cBCD products, we next applied several steps to correct for variations and features in the image backgrounds. We corrected each image for column-pulldown effects associated with bright sources using custom software (“fixpull-down.pro”)<sup>4</sup>. We next constructed a median frame from all cBCDs in a single AOR, clipping outliers. In this way we make a sky frame from all images in an AOR. We subtracted this image from each individual cBCD to eliminate structure and residuals. We then removed additional striping in the backgrounds by averaging over five columns in each image (clipping for objects), and subtracted this from each column. We also excluded the first frame from each series of exposures in a given AOR sequence to remove the any variable instrument bias level associated with idiosyncrasies of the post-


cryogenic IRAC electronics (the “first frame effect”<sup>5</sup>).

Our inspection of preliminary reductions showed some instances where persistence from bright stars produced spurious sources in the final mosaics. This occurs even though the data-reduction pipeline flags for image persistence (and we set persistence-flagged pixels as “fatal” during the mosaic stage, see below), and suspect this occurs because the pipeline flags only persistence from the brightest objects (and fainter objects, which still cause persistence, are missed). Figure 2 shows an example of the persistence caused by a bright star (HD 9670,  $V = 6.9$  mag) in a consecutive series of cBCDs from one of the SHELA AORs. Following our observing strategy, the star is dithered to 3 different positions on the detector, before the IRAC array is stepped to a new location on the sky. The persistence from the star is evident in several subsequent exposures. The persistence fades with an exponential timescale (as expected for trapped electron decay rates) and is mainly a problem in the  $3.6 \mu\text{m}$  images (it is nearly negligible in the  $4.5 \mu\text{m}$  images, see footnote 5).

To correct for the strongest persistence residuals, one of us

<sup>4</sup> <http://irsa.ipac.caltech.edu/data/SPITZER/docs/dataanalysisstools/tools/contributed>

<sup>5</sup> See the IRAC Instrument Handbook (2015, version 2.1), available at: <http://irsa.ipac.caltech.edu/data/SPITZER/docs/irac/iracinstrumenthandbook>



**Figure 5.** Combined, three-epoch exposure-time map of the SHELA IRAC Channel 1 ( $3.6 \mu\text{m}$ ) data.

(CL) inspected visually each channel 1 cBCD sequentially in the order they were observed, identifying persistence events. We then flagged those pixels with persistence using the locations of the bright objects in the previous cBCDs in the observing sequence. We combined these flag maps with the mask files produced by the SSC pipeline and included the masked pixels as fatal bits in the mosaicking steps. Even so, we have likely not accounted for all possible persistence in the images. Persistence can manifest as “sources” that vary in the time domain between observations in different epochs, and users of the catalogs (especially for time-domain studies or sources detected in a single channel of an observing epoch) should be wary that some time-variable sources may be a result of faint persistence missed by our inspection of the images.

### 3.2. Image Mosaics

We used a combination of the MOPEX software (v18.5.4) provided by the SSC<sup>6</sup> and SWarp (v2.19.1 Bertin et al. 2002) to produce mosaics of the IRAC data. Our choice to use SWarp is a result of the fact that the memory limitations of

MOPEX are too stringent for a dataset with the size of the SHELA data volume. We first produced a mosaic for each AOR separately using MOPEX. MOPEX includes full propagation of errors for each pixel and masks pixels set to fatal bit patterns (including pixels we estimate to contain persistence, see above).

We next used SWarp to mosaic the output from MOPEX for each AOR into images covering the full SHELA field. We employed a background subtraction with `BACK_SIZE=128` and `BACK_FILTERSIZE=3` within SWarp to account for (small) offsets in the backgrounds between AORs. We combined AORs using a weighted average (`COMBINE_TYPE=WEIGHTED`) from the exposure-time maps for each AOR, and we resampled the images to a common field center and pixel scale of  $0''.8 \text{ pixel}^{-1}$ . We produced full mosaics of all the data at  $3.6 \mu\text{m}$  and  $4.5 \mu\text{m}$ . We also produced mosaics in each channel in each of the 3 observing epochs separately. Figures 3 and 4 show the combined, three-epoch mosaics at  $3.6$  and  $4.5 \mu\text{m}$ , respectively.

We combined the weight maps using SWarp. Figures 5 and 6 show the weight map coverage for the full mosaic at  $3.6$  and  $4.5 \mu\text{m}$ , respectively. The values in the weight map correspond to the number of IRAC exposures for each pixel on the

<sup>6</sup> <http://irsa.ipac.caltech.edu/data/SPITZER/docs/dataanalysis/tools>






Figure 6. Same as Figure 5, for the SHELA IRAC Channel 2 ( $4.5 \mu\text{m}$ ) data.




Figure 7. The cumulative distribution of area with exposure times  $\geq t_{\text{exp}}$ . The thin lines show the distribution only for the first epoch IRAC data, and the thick lines show the distribution for the combined three epoch data. The solid lines show the  $3.6 \mu\text{m}$  distribution and the dotted line shows the  $4.5 \mu\text{m}$  distribution.

sky, and the weight map is therefore proportional to the effective exposure time, where the exposure time for each image is  $t_{\text{eff}} = 23.6 \text{ s} \times W$ , where  $W$  is the value of the weight map. Figure 7 shows the distribution of area covered to a given exposure time in the  $3.6 \mu\text{m}$  and  $4.5 \mu\text{m}$  full mosaics compared to the coverage from epoch 1 only. A single epoch covers each region of a  $26 \text{ deg}^2$  area with three pointings from the dither pattern ( $W = 3$ ), for an effective exposure time,  $t_{\text{eff}} \simeq 71 \text{ s}$ . The full mosaic covers an area of  $30 \text{ deg}^2$  with  $W = 6$  pointings for  $t_{\text{eff}} = 142 \text{ s}$  depth,  $24.2 \text{ deg}^2$  with at least 200 s depth ( $W = 8.5$ ), and it covers  $22.4 \text{ deg}^2$  with  $W = 9$  pointings for  $t_{\text{eff}} = 212 \text{ s}$  depth (the nominal, full 3 epoch depth).

### 3.3. Astrometric Quality

In preliminary versions of the SHELA IRAC mosaics, we identified small astrometric offsets between cBCDs from different AORs. On subsequent re-reductions, we correct for these inter-AOR shifts using multiple tests.

We computed coarse astrometric offsets by cross-correlating the positions of objects in each cBCD with those detected in the SDSS DR7 (Abazajian et al. 2009) catalog and updating the image headers. The astrometric offsets were mostly small, with shifts of up to  $\approx 0''.2$  in both R.A. and

Decl. We then combined all cBCDs from all mosaics from each epoch, and we again checked the absolute astrometry of each epoch of the SHELA data, using the newer SDSS DR7 as a reference frame. We corrected for the remaining (small) relative shifts between each epoch. Compared to SDSS DR7 the offsets of the 3.6 micron images were:  $\Delta\alpha = \alpha_{\text{SHELA}} - \alpha_{\text{DR7}} = -140, +180, -140$  mas, and  $\Delta\delta = \delta_{\text{SHELA}} - \delta_{\text{DR7}} = +60, -80, +50$  mas, for epochs 1, 2, and 3, respectively. The offsets to the 4.5 micron images were slightly different:  $\Delta\alpha = -134, +140, -130$  mas; and  $\Delta\delta = -140, +110, -150$  mas, for epochs 1, 2, and 3, respectively. The origin of the offsets is unclear, but may be related to the errors measured in the positions of the stars in the *Spitzer* star trackers for the different spacecraft orientations (where the errors may be a combination of uncertainties in the proper motions in the guide star catalog, proper motions of stars in the 2MASS catalog used for the pointing refinement step of the IRAC pipeline, and intrapixel sensitivity variations that add noise to the measured star positions). The orientation of epochs 1 and 3 were approximately the same, while the spacecraft orientation for epoch 2 was different by approximately 180 degrees, and indeed, the largest offsets were between epoch 1 and 2 and epochs 2 and 3 (see above). We corrected for these astrometric offsets between each epoch before combining the data into the final mosaics. Our tests showed that correcting for astrometric shifts for each epoch improved the image quality of point sources in the final mosaic.

We remosaicked all the data using the improved astrometric corrections. As a result, the final astrometric solutions are very good compared to SDSS DR7. Figure 8 shows that offsets between SHELA and DR7 are indeed very small,  $\Delta\alpha = \alpha_{\text{SHELA}} - \alpha_{\text{DR7}} = -14$  mas and  $\Delta\delta = \delta_{\text{SHELA}} - \delta_{\text{DR7}} = 7$  mas. For comparison, the scatter is  $\sigma(\alpha) = 180$  mas and  $\sigma(\delta) = 160$  mas in each dimension, respectively. The scatter is comparable to the quoted uncertainty in the SDSS DR7 astrometric solution (Abazajian et al. 2009). We have also rechecked the astrometry between SHELA and the newer SDSS DR9, which are similar:  $\Delta\alpha = -23$  mas and  $\Delta\delta = 18$  mas, with corresponding scatter  $\sigma(\alpha) = 160$  mas and  $\sigma(\delta) = 150$  mas. The slight increase in the offset between DR7 and DR9 is well within the uncertainty in the absolute astrometric calibration of SDSS (Ivezic et al. 2007). The offsets are also small between SDSS DR7 and the IRAC images from each individual epoch.

We also compared the astrometry between SHELA and the 2MASS all-sky point-source catalog (Skrutskie et al. 2006). There are very small shifts of  $\Delta\alpha = -8$  mas and  $\Delta\delta = -30$  mas, with scatter  $\sigma(\alpha) = 270$  mas and  $\sigma(\delta) = 260$  mas in each dimension. This is larger than the typical positional uncertainty for  $K_s < 14$  mag sources ( $\lesssim 100$  mas, Skrutskie et al. 2006), but we have made no correction for proper motion of stars, and the accuracy is consistent with that reported in Sanders et al. (2007, who state an accuracy of  $\sim 200$  mas for their *Spitzer* IRAC data).

There are larger shifts between the SHELA IRAC astrometry and the astrometry of point sources in the AllWISE catalog (Cutri et al. 2013). Figure 8 shows that the offsets are  $\Delta\alpha = -53$  mas and  $\Delta\delta = 69$  mas, with scatter of  $\sigma(\alpha) = 190$  mas and  $\sigma(\delta) = 200$  mas in each dimension. The scatter is consistent with the astrometric uncertainty of the AllWISE catalogs (Cutri et al. 2013), but the larger offsets in the astrometry (approaching a tenth of an arcsecond) may be non-negligible for some applications.

### 3.4. Point Response Functions

For a variety of quality tests of the data and catalogs, it is useful to have an empirical point response function (PRF) for the 3.6 and 4.5  $\mu\text{m}$  images. In each image, we identified point sources from the AllWISE catalog brighter than  $W1 < 15$  Vega mag with flag value  $e_x = 0$  within the SHELA footprint. We kept objects only in the magnitude range  $14 < W1 < 15$  Vega mag as these have high signal-to-noise, are well away from being saturated, and because brighter objects are weighted more in the construction of the PRF. We constructed average PRFs using the routines provided in IDLPhot<sup>7</sup>, based on the DAOPhot software (Stetson 1987). Figure 9 shows the PRFs for the 3.6 and 4.5  $\mu\text{m}$  data. We use the empirical PRF for tests of object photometric accuracy and completeness in § 4.1 and § 4.2 below. We measure a full-width at half maximum (FWHM) for  $1''.97$  and  $1''.99$  from Gaussian fits to the 3.6  $\mu\text{m}$  and 4.5  $\mu\text{m}$  PRFs, respectively. These agree with the expected values at the native IRAC pixel scale for the IRAC channel 1 and 2 detectors during the warm mission.<sup>8</sup>

We measured a curve-of-growth of the PRFs using circular apertures and compared those to the flux measured with the fiducial IRAC aperture (radius  $R = 12''$ ) used to derive the IRAC flux calibration (see the IRAC Instrument Handbook, link in footnote 5). The curve of growth provides an estimate of the amount of light lost outside the photometric aperture. For large apertures ( $R > 2''$ ) these corrections are identical to the ones we adopt below (§ 3.5), but they differ at the 0.05–0.10 mag level for apertures  $R < 1-2''$ .

### 3.5. Photometric Aperture Corrections

Because we used the SExtractor software package for object photometry in our SHELA catalog (§ 4, below), we found it advantageous to derive aperture corrections for point-source photometry from the images themselves using the same photometric software package. We used the same bright ( $W1 < 15$  Vega mag) objects selected from the AllWISE catalogs used for the construction of the PRF (see § 3.4). We subsequently photometered those objects in the IRAC images using SExtractor with the same parameters as the source catalog (see § 4 and Table 3), using the AllWISE point sources as an associated list with a search radius of 5 pixels ( $4''$ ). SExtractor photometered these sources with circular apertures ranging in radius from  $1''$  to  $12''$ , where the  $12''$ -radius aperture “defines” the total aperture (see § 3.4, above).


Figures 10 and 11 compare the photometric magnitudes measured in different-sized apertures to the total,  $R = 12''$  aperture for the IRAC 3.6 and 4.5  $\mu\text{m}$  data, respectively. As expected, there are offsets owing to light lost outside the smaller apertures. The offsets are constant for sources,  $\gtrsim 13.5$  mag in IRAC. We measure aperture corrections based on the median  $m(< R) - m(< 12'')$  magnitude for stars with magnitudes between 13.5 and 16 (AB) mag. These aperture corrections are shown as long-dashed, red lines in each panel in Figures 10 and 11. For brighter sources, the offset is a function of source magnitude, and we caution using photometry measured in small apertures for sources brighter than  $\lesssim 14$  mag as the aperture corrections are unreliable.

Figure 12 shows the aperture corrections for the IRAC 3.6


<sup>7</sup> <http://idlastro.gsfc.nasa.gov/ftp/pro/idlphot>

<sup>8</sup> Data taken with IRAC during the warm *Spitzer* mission have measured FWHMs for the PRFs  $\approx 15\%$  larger than for data taken during the cold *Spitzer* mission, see the IRAC instrument handbook, link in footnote 5.





**Figure 8.** Astrometric comparison for point sources between the SHELA/IRAC and SDSS DR7 (left panel) and between SHELA/IRAC and AllWISE (right panel). In each panel, the shading is proportional to the density of objects in each region of the plot. The red, dashed lines show the mean offset. The ellipses contain 68% of the sources. There are negligible offsets between SDSS DR7 and SHELA (by construction), and the scatter is consistent with the SDSS astrometric accuracy of  $0''.2$  (Abazajian et al. 2009). In contrast, the AllWISE astronomy has a non-negligible offset of  $\approx 60$ – $70$  mas in both R.A. and Decl., with a scatter of  $0.2''$  in each dimension.



**Figure 9.** Empirical PRFs for the SHELA IRAC 3.6 and 4.5  $\mu\text{m}$  data. The PRFs are constructed by combining the IRAC fluxes for point sources from the AllWISE catalog with magnitudes  $14 < W1 < 15$  Vega mag.

**Table 2**  
Aperture Corrections for IRAC Data

Radius, $R$ (pix) (1)	Radius, $R$ (arcsec) (2)	[3.6](< $R$ ) - [3.6](< 12) (mag) (3)	[4.5](< $R$ ) - [4.5](< 12) (mag) (4)
2.500	2.0	0.326	0.353
3.125	2.5	0.213	0.226
3.750	3.0	0.157	0.159
5.000	4.0	0.107	0.112
6.250	5.0	0.074	0.085
7.500	6.0	0.053	0.058

**Note.** — The aperture correction is the difference between the magnitude measured in a circular aperture of radius  $R$  and the magnitude measured in a circular aperture of radius  $12''$ .

and 4.5  $\mu\text{m}$  data measured for point sources with magnitude between 13.5 and 16 mag in circular apertures of radius  $R$  and a total magnitude measured in the total,  $R = 12''$  aperture. Table 2 lists the aperture correction for each band for the apertures included in the catalogs here. These aperture corrections are consistent with those of the IRAC Instrument Handbook (see footnote 5) and those derived in the literature (e.g., Ashby et al. 2009, 2013b) with differences at the  $< 0.05$  mag level. These differences likely depend on the method of photometry. We advocate the use of the aperture corrections derived here as they use the same photometric parameters as the

source catalog. These corrections are accurate to better than 0.03 mag based on our comparison of the IRAC photometry to flux measurements from AllWISE at W1 (3.4  $\mu\text{m}$ ) and W2 (4.6  $\mu\text{m}$ ) in § 3.6, below.


### 3.6. Photometric Quality: Comparison between SHELA IRAC and AllWISE

A test of the photometric accuracy of the SHELA IRAC data is possible by comparing the IRAC photometry to that measured by WISE at 3.4  $\mu\text{m}$  (W1) and 4.6  $\mu\text{m}$  (W2). For this test, we use the same point sources selected from the AllWISE catalog matched to the IRAC data discussed in § 3.5, above. Figure 13 shows the magnitude difference between the IRAC [3.6] and WISE W1 photometry and between IRAC [4.5] and WISE W2 photometry. The AllWISE W1 and W2 catalogs have  $5\sigma$  sensitivity limits  $\gtrsim 16$  mag (Vega, Cutri et al. 2013) so our comparison is for stars well above this limit and biases should be minimal. The difference between [3.6]–W1 is negligible. The figure shows this distribution, where a Gaussian fit to it gives a mean  $\mu = 0.001$  mag and standard deviation,  $\sigma = 0.023$  mag.

Figure 13 shows the distribution of [4.5]–W2, where a Gaussian fit gives mean  $\mu = -0.028$  mag with  $\sigma = 0.021$  mag. While the offset is small ( $-0.028$  mag), its origin is unclear. The offset is  $1.4\sigma$  times the scatter, and corresponds to a flux ratio of 1.8%. Jarrett et al. (2011) find similar offsets and conclude these are consistent with the differences in the IRAC and WISE bandpasses (the “relative system response” curves), and the uncertainty in the calibration. A similar offset is found by Cutri et al. (2013) between W2–[4.5]<sup>9</sup>. This offset persists regardless of the size of the photometric aperture (once corrected to total), so a systematic offset in the aperture correction seems unlikely. It therefore seems to be consistent with differences in the WISE W2 and IRAC [4.5] spectral response curves.

We furthermore considered (and rejected) the possibility that there are additional color terms between IRAC and WISE photometry. Figure 14 shows the expected color between the IRAC and WISE bands for different stellar types, using models from Kurucz (1993) over a range of luminosity class and

<sup>9</sup> see [http://wise2.ipac.caltech.edu/docs/release/allwise/expsup/sec2\\_3a.html](http://wise2.ipac.caltech.edu/docs/release/allwise/expsup/sec2_3a.html)



**Figure 10.** Estimate of light lost outside circular apertures of varying radius in the SHELA IRAC 3.6  $\mu\text{m}$  image. Each panel shows the difference between a total magnitude (defined in a  $12''$ -radius aperture) and the magnitude measured in a smaller circular aperture of radius  $R$  for point sources from the AllWISE catalog. Each panel shows a different aperture radius  $R$ , given above each panel. The dashed, thick red line in each panel shows the median difference for stars between  $13.5 < [3.6] < 16$  mag, used to derive the aperture correction.

spectral type. The IRAC–WISE colors are zero (relative to Vega) for early-type (i.e., Vega-analogs) main-sequence stars. This is expected as these stars are used for the calibration of the instruments.<sup>10</sup> For example, because the WISE bands are broader in wavelength than the IRAC bands, they are more sensitive to the molecular absorption in later-type stellar atmospheres (e.g.,  $\text{H}_2\text{O}$ ,  $\text{HCN}$ ,  $\text{C}_2\text{H}_2$ ,  $\text{CO}$ ), especially in the giants, supergiants, and AGB stars that may contribute substantially to the stellar counts in the SHELA data. Figure 14 shows that the atmospheres of late-type supergiants show color offsets of  $[4.5] - W2 \simeq -0.02$  mag, consistent with the offset observed in the data and other comparisons in the literature (e.g., Jarrett et al. 2011). However, these same stars have about the same offset in  $[3.6] - W1$  (although as a result of different absorption features,  $\text{H}_2\text{O}$  and  $\text{C}_2\text{H}_2 + \text{HCN}$  bands at  $< 3.8 \mu\text{m}$  versus  $\text{CO}$  band at  $5.0 \mu\text{m}$ ; see Matsuura et al. 2005, 2014), which calls into doubt this interpretation. Therefore, while color terms likely can explain the *scatter* in the IRAC–AllWISE colors (Figure 13), they likely are not the cause of the systematic offset between the IRAC 4.5  $\mu\text{m}$  and AllWISE

<sup>10</sup> Because the IRAC and WISE filters have different widths, there are color terms for magnitudes measured in these filters when converted to AB units even though the colors for Vega-type stars are  $\approx 0.0$  mag. This is evident from the quoted AB-to-Vega conversions. The AB-to-Vega conversions for WISE are  $W1_{\text{AB}} - W1_{\text{Vega}} = 2.699$  mag and  $W2_{\text{AB}} - W2_{\text{Vega}} = 3.338$  (Jarrett et al. 2011). Comparing to the conversions for IRAC (see § 1), this implies a IRAC – WISE color of  $\approx 0.1$  mag color for Vega-like stars on the AB magnitude system (see also, Richards et al. 2015).

4.6  $\mu\text{m}$  photometry discussed in the previous paragraph.

Regardless of its origin, the offset is small, and is within the uncertainty of the absolute IRAC calibration (Reach et al. 2005). Jarrett et al. (2011) argue the offset likely results from a combination of absolute calibrations, aperture corrections, and/or color corrections. The photometry is sufficiently accurate for most science applications, although those requiring better than 2% absolute photometric accuracy should be aware of this systematic.

#### 4. SHELA IRAC CATALOGS

We used Source Extractor (SExtractor v. 2.19.5; Bertin & Arnouts 1996) to detect and to photometer sources in the IRAC images. To detect sources, we constructed a detection image as the weighted sum of the 3.6 and 4.5  $\mu\text{m}$  images. The detection image,  $D$ , is then,

$$D = \frac{W_1 \times I_1 + W_2 \times I_2}{W_1 + W_2} \quad (1)$$

where  $W_1$  and  $W_2$  are the weight maps (proportional to the exposure time) for the IRAC channel 1 and 2 images, respectively, and  $I_1$  and  $I_2$  are the science (flux) images for channel 1 and 2, respectively. We then ran SExtractor in “double image mode” using the detection image and science images with the parameters listed in Table 3. Tables 7 and 8 provide all the information from the full-mosaic catalogs.

We also constructed catalogs for the mosaics from each of the three observing epochs individually. For each epoch,






Figure 11. Same as Figure 10 for the SHELA IRAC 4.5  $\mu\text{m}$  image.




Figure 12. Aperture corrections for the IRAC images. The plot shows the difference between the IRAC photometric magnitude measured for point sources with magnitude between 13.5 and 16 mag in circular apertures of radius  $R$  and a total magnitude measured in an  $R = 12''$  aperture. The aperture corrections are measured in apertures with discrete radii, as indicated by the solid black squares, and interpolated linearly between those points. The solid (dashed) line shows the 3.6  $\mu\text{m}$  (4.5  $\mu\text{m}$ ) data, as labeled in the plot legend.

we used the detection for the combined images (see eq. 1 above). In this way, sources detected in the combined epoch, 3.6+4.5  $\mu\text{m}$  image are photometered in each image from each epoch. We use the identical SExtractor parameters as for the full-mosaic catalogs (Table 3). Tables 9–11 provide the photometry from the individual epoch data.

Table 3  
SHELA SExtractor Parameter Settings

SExtractor Parameter (1)	Value (2)
DETECT_MINAREA	3 pixels
DETECT_THRESH	1.5
ANALYSIS_THRESH	1.5
FILTER_NAME	gauss_2.0_5x5 <sup>a</sup>
WEIGHT_TYPE	MAP_WEIGHT
DEBLEND_NTHRESH	64
DEBLEND_MINCONT	0.0005
MAG_ZEROPOINT	20.9555 <sup>b</sup>
PIXEL_SCALE	0.80 arcsec
BACK_TYPE	AUTO
BACK_SIZE	256 pixels
SEEING_FWHM	1.7 arcsec


**Note.** — SExtractor was run using the weighted sum of the 3.6 and 4.5  $\mu\text{m}$  images for detection, and using the images separately for photometry. All SExtractor parameters are identical for both images. All other SExtractor parameters are set to the program defaults (for SExtractor v.2.19.5).

<sup>a</sup> This is a Gaussian kernel with  $\sigma=2$  pixels and size  $5 \times 5$  pixel<sup>2</sup> used to filter the image for source detection.


<sup>b</sup> The AB magnitude zeropoint for the images, converting from the *Spitzer* default of MJy sr<sup>-1</sup> to  $\mu\text{Jy pixel}^{-1}$  at the  $0''.8$  pixel<sup>-1</sup> scale.

#### 4.1. Completeness Simulations

We performed simulations to estimate the completeness in the SHELA IRAC catalogs following the method in Papovich et al. (2015). We inserted fake point sources into the 3.6 and 4.5 images using the empirical PRFs derived above (§ 3.4). We inserted each fake source at the same  $(\alpha, \delta)$  lo-



**Figure 13.** Comparison of magnitudes measured for point sources in the AllWISE catalog at  $3.4 \mu\text{m}$  ( $W1$ ) and  $4.6 \mu\text{m}$  ( $W2$ ) compared with those in the SHELA IRAC  $3.6$  and  $4.5 \mu\text{m}$  data. Note that in this plot all magnitudes are relative to Vega-type stars. In both plots the gray shading shows all point sources, where darker regions correspond to a higher density of points falling in that region. The red points show a random subsample of the data. In each plot, the right panel shows the distribution of the magnitude difference between the IRAC and WISE image. The left plot compares the IRAC  $3.6 \mu\text{m}$  and WISE  $W1$  data. The photometric offset is negligible. The right plot compares the IRAC  $4.5 \mu\text{m}$  and WISE  $W2$  data. There is a small offset,  $-0.028 \text{ mag}$ . Stars brighter than  $\lesssim 10$  ( $11$ ) Vega mag appear saturated in the IRAC  $3.6$  ( $4.5$ )  $\mu\text{m}$  data.



**Figure 14.** Expected differences between the IRAC and WISE photometry due to color variations in stars of different spectral type. The top panel compares the transmittance of the IRAC  $[3.6]$  and  $[4.5]$  filters to the WISE  $W1$  and  $W2$  filters. The plot shows synthesized colors between IRAC  $3.6 \mu\text{m}$  and WISE  $W1$ , and between IRAC  $4.5 \mu\text{m}$  and WISE  $W2$ . The data points correspond to Kurucz (1993) models for dwarfs (luminosity class V), giants (class III), and supergiants (class I) over a range of effective temperature (spectral type). Because the WISE instrumental filters are wider, they can include bandhead absorption features in late-type stars, affecting the IRAC–WISE color up to  $0.03 \text{ mag}$ .

cation in the  $3.6$  and  $4.5 \mu\text{m}$  images, where the source has the same total brightness (AB magnitude) in each channel. Fake sources have magnitudes chosen from a wide distribution ( $17$ – $24$ th magnitude), and the sources are located anywhere in the images. In this way fake sources may fall within the isophote of real objects in the image, and therefore our completeness simulations include the effects from blended objects. We reconstruct the detection image as the weighted sum of the  $3.6$  and  $4.5 \mu\text{m}$  images and rerun SExtractor. This


**Table 4**  
Completeness and Error Estimates for SHELA IRAC data

AB mag (1)	Raw Completeness (2)	Completeness (3)	$\sigma_{3.6}$ (4)	$\sigma_{4.5}$ (5)
18.0	0.98	0.98	0.02	0.02
18.5	0.97	0.97	0.02	0.02
19.0	0.98	0.98	0.02	0.02
19.5	0.97	0.97	0.03	0.03
20.0	0.95	0.95	0.05	0.05
20.5	0.94	0.94	0.07	0.07
21.0	0.90	0.90	0.11	0.12
21.5	0.86	0.86	0.17	0.17
22.0	0.79	0.79	0.25	0.25
22.5	0.57	0.56	0.37	0.38
23.0	0.20	0.19	0.49	0.51
23.5	0.05	0.04	0.76	0.77
24.0	0.01	0.00	0.90	0.87

**Note.** — (1) Magnitude bin, (2) ratio of the number of recovered fake sources to the total number of fake sources in this magnitude bin, (3) completeness corrected for “false positives”, fake sources recovered in the detection image even when no fake sources were added, (4) estimate of the photometric uncertainty for point sources in  $[3.6]$ , (5) estimate of the photometric uncertainty for point sources in  $[4.5]$ . The photometric uncertainty estimates are the standard deviation between the input magnitudes and the measured magnitudes (measured in  $2''$ -radius apertures, corrected to total using the values in Table 2).

latter step is computationally expensive given the size of the images (see above; Equation 1). We repeated the simulation only 15 times, where we inserted into each simulated image 10,000 fake sources ( $\approx 0.4\%$  the total number of real sources). In this way we sample the full range of source magnitude using the minimum investment of resources.

We computed the completeness as the ratio of the number of recovered (detected) fake sources to the number of input fake sources in bins of source magnitude. Figure 15 shows the completeness, where the 50% (80%) completeness limit is 22.6 (22.0) AB mag. Table 4 gives these as the “raw” completeness as a function of source magnitude in the detection image. We also added to the completeness a correction for “false positives”, sources at the location of the fake that are “recovered” even when no sources are added to the image. Table 4 gives these as the “completeness”. As illustrated in Figure 16 the difference between the raw completeness and



**Figure 15.** Completeness for point-sources in the SHELA data as a function of the input source magnitude. The plot shows the recovery fraction as a function of magnitude for simulated point sources, which are added to both the 3.6 and 4.5  $\mu\text{m}$  images with the same AB magnitude. The solid-line histogram shows the raw completeness fraction. The dashed-line histogram shows the completeness corrected for “false positives” (sources that are “recovered” in the image, in which no simulated sources are added). The solid, dashed, and dot-dashed horizontal lines show 100%, 80%, and 50% completeness.

the completeness corrected for false positives is small, accounting for only 1% of recovered sources down to the 50% completeness limit.

#### 4.2. Error Estimates

We estimate uncertainties for sources in the IRAC catalogs using two methods. We first used the simulations from § 4.1 to estimate the uncertainty for point sources of a given magnitude. In each of the IRAC 3.6 and 4.5  $\mu\text{m}$  images, we computed the difference between the input (“true”) magnitude and measured magnitude from SExtractor in  $R = 2''$  radii (corrected to total using the aperture corrections in Table 2). Figure 16 shows the median and inter-68-percentile of the distributions of these differences as a function of [3.6] and [4.5] magnitude. The mean offset is near zero down to  $\approx 22.2$  mag (below the 80% completeness limit). In each bin of magnitude, we compute the ratio  $\sigma/F_\nu$  (the inverse of the signal-to-noise ratio, SNR) as a measure of the relative error for sources of that magnitude. The solid black line of Figure 17 shows this ratio as a function of [3.6] and [4.5]. This yields a limiting SNR=5 at 22.0 AB mag, or a  $1\sigma$  limit of 1.1  $\mu\text{Jy}$  for both [3.6] and [4.5].

This  $1\sigma$  flux-density limit is consistent with estimates from the *Spitzer* SENS-PET for fields with higher background. SENS-PET gives for the “warm” *Spitzer* mission a  $1\sigma$  limit for point sources of 0.9–1.2  $\mu\text{Jy}$  and 1.1–1.6  $\mu\text{Jy}$  for 3.6 and 4.5  $\mu\text{m}$  observations for “medium” and “high” background (where as noted above, the medium and high backgrounds in SENS-PET assume a sightline with latitude  $\beta = 40^\circ$  and  $0^\circ$  from the Ecliptic); therefore the values we derive are reasonable.

Figure 17 shows a “kink” in the error estimated from the simulations for both [3.6] and [4.5] below  $\gtrsim 22.5$  mag (black, solid-lined curves in each panel of the figure). This is likely a bias owing to incompleteness of recovered sources. The 50 and 80% completeness limits are 22.6 and 22.0 mag, re-

spectively. Figure 16 shows that in this magnitude range, the median difference between recovered and input photometry is biased to positive values because fainter sources are missed in the catalog. Therefore, at these magnitudes, the distribution is clipped, and the inter-68 percentile range is biased smaller. This means the errors estimated from the simulations underestimate the true photometric uncertainty for sources with magnitudes below about the 80% completeness limit. Partly for this reason we will adopt the alternative method to estimate errors, described in the rest of this subsection below.

Second, we derived error estimates from the noise in the images in apertures of increasing linearized number of pixels,  $N$ , where  $N = \sqrt{A}$  and  $A$  is the area of the photometric aperture. The flux uncertainty within an aperture has a contribution from photon statistics. The theoretical uncertainty in an aperture with  $N$  pixels would then scale as  $\sigma_N = \sigma_1 \times N$ , where  $\sigma_1$  is the standard deviation of background pixels. This relation assumes pixels are independent (uncorrelated), and in practice this is imperfect as the image reduction, data processing, and mosaicking introduce pixel-to-pixel correlations, in addition to uncertainties in background subtraction, undetected objects, and extended emission in the wings of bright objects. The limiting case of perfect correlations between pixels implies that the uncertainty in an aperture of  $N$  linearized pixels should scale as  $\sigma_N = \sigma_1 \times N^2$  (Quadri et al. 2007). Generalizing, we expect the uncertainty to scale with  $N^\beta$  with  $1 < \beta < 2$ , between the limiting cases of uncorrelated pixels and perfectly correlated pixels (see also Labbé et al. 2003; Gawiser et al. 2006; Blanc et al. 2008; Whitaker et al. 2011).

We estimated the noise as a function of linearized pixels by measuring the sky counts in circular apertures of varying size in  $\approx 5000$  randomly placed regions in the SHELA IRAC images, ensuring that apertures do not overlap, and excluding regions containing objects. We then computed the standard deviation of the distribution of aperture fluxes from the normalized median absolute deviation,  $\sigma_{\text{nmad}}$  (Beers et al. 1990), as an estimate for  $\sigma_N$  for each aperture with  $N$  linearized pixels. Figure 18 shows the measured relation of  $\sigma_N$  as a function of  $N$  for the 3-epoch IRAC [3.6] and [4.5] images.


Following the suggestion in Labbé et al. (2003), we fit a parameterized function to estimate the noise in an arbitrary aperture of linear size  $N$ ,

$$\sigma_N = \sigma_1 (\alpha N^\beta + \gamma N^\delta) \quad (2)$$


where  $\sigma_1$  is the pixel-to-pixel standard deviation in the sky background, and  $\alpha$ ,  $\beta$ ,  $\gamma$ , and  $\delta$  are free parameters. We required that  $\alpha$  and  $\gamma$  be non-negative, that  $1 < \beta < 2$ , and we placed no restrictions on  $\delta$ . In this way the first term of Equation 2 represents the expected noise for partially correlated pixels. The second term includes an additional correction that better reproduces the noise in large apertures (see also, Labbé et al. 2003). Table 5 lists the parameters for the fits in Equation 2 for the combined, 3 epoch IRAC 3.6 and 4.5  $\mu\text{m}$  data. The table also includes fits for the individual IRAC epoch data (where our tests showed each individual epoch had noise properties consistent with being the same, so we combined the random apertures of all individual images for a single fit).

The product of  $\sigma_1 \alpha$  reflects the pixel-to-pixel rms, and these decrease roughly with the square-root of the exposure time such that the value of  $\sigma_1$  for the 3-epoch combined image is roughly  $\sqrt{3}$  lower than that for an individual epoch. The fitted values for the slope in the first term,  $\beta \sim 1.3-1.4$ , are consis-






**Figure 16.** Comparison between the “true” (input) magnitude for simulated sources and the measured magnitude as a function of sources magnitude. The left panel shows the results for  $3.6 \mu\text{m}$ , the right panel shows the results for  $4.5 \mu\text{m}$ .



**Figure 17.** Estimates of uncertainties for the SHELA IRAC data. The left panel shows the results for the IRAC  $3.6 \mu\text{m}$  data. The right panel shows the results for the  $4.5 \mu\text{m}$  data. In each panel, the solid-line curve shows the estimated uncertainty measured from a comparison of the recovered magnitudes to the input magnitudes for fake sources added to the images. The points connected by the dashed-line curve show the estimates derived from  $\sigma_N$  for  $2''$ -radii apertures, scaled to total magnitudes. The horizontal lines show the equivalent magnitude uncertainty for a source with  $\text{SNR}=5, 3$ , and  $2$ , as labeled.

**Table 5**  
Coefficients for Error Estimates using  $\sigma_N$

Channel (1)	Epoch (2)	$\sigma_1/\mu\text{Jy}$ (3)	$\alpha$ (4)	$\beta$ (5)	$\gamma$ (6)	$\delta$ (7)
$3.6 \mu\text{m}$	Combined 3 Epochs	0.106	0.959	1.33	$8.80 \times 10^{-4}$	3.53
	Single Epoch	0.178	0.944	1.37	$1.98 \times 10^{-3}$	3.76
$4.5 \mu\text{m}$	Combined 3 Epochs	0.103	0.981	1.34	$3.49 \times 10^{-4}$	4.55
	Single Epoch	0.169	0.893	1.39	$1.95 \times 10^{-6}$	5.17



**Figure 18.** Scaling relation between the measured noise in the SHELA IRAC images and the number of linearized pixels,  $N = \sqrt{A}$  (where  $A$  is the area of the photometric aperture). Both panels show the measured noise,  $\sigma_N$ , in each aperture  $N$ . The top panel shows the  $3.6 \mu\text{m}$  data and the bottom panel shows the  $4.5 \mu\text{m}$  data. In each panel, the bottom-most dashed line shows the theoretical relation assuming uncorrelated pixels in the Gaussian limit,  $\sigma_N \sim N$ . The top-most dashed line shows the relation for perfectly correlated pixels ( $\sigma_N \sim N^2$ , Quadri et al. 2007). The red, short-dashed line shows the parameterized fit to the data, which we use to define the flux uncertainties measured in different-sized apertures.

tent with partially correlated pixels, as found in other imaging surveys (e.g., Gawiser et al. 2006; Quadri et al. 2007). The values for  $\gamma$  are relatively small (the ratio of the coefficients is  $\alpha/\gamma \sim 5 \times 10^2 - 5 \times 10^5$ ), implying there is a small, but increasing correction to the noise model for apertures with larger numbers of pixels.

The red points in Figure 17 show the magnitude uncertainty calculated for  $2''$ -radius apertures (scaled up to the total aperture) as a function of [3.6] and [4.5]. There is generally good agreement between the estimated uncertainties from  $\sigma_N$  and those from the simulations described above. For objects with  $< 22$  AB mag, there is a slight offset, where the estimates from  $\sigma_N$  are lower at about the 0.02 mag level compared to the estimates from the completeness simulations. This could arise from several effects, including the fact that the completeness simulations allow fake objects to fall on image regions that contain other (real) galaxies. As this will tend to increase the average difference between the input and recovered magnitude, an offset is not unexpected. While we do not include this additional error in our catalogs, the magnitude uncertainties in our catalog may require an additional 0.02 mag sys-

tematic uncertainty to account for this effect.

For the IRAC catalogs, we computed errors using Equation 2 for the linearized number of pixels  $N = \sqrt{A}$  where  $A$  is the total area of the aperture used to measure the object, scaled up to the total aperture. We add these errors in quadrature with an additional error,  $\sqrt{\text{ADU}/G}$ , to account for poisson uncertainties owing to the counts associated with each object, where ADU is the astronomical data unit, and the gain is  $G = 3.7 \text{ e}^-/\text{ADU}$  for post-cryogenic *Spitzer*/IRAC observations. We opted to use these uncertainty estimates as they can be scaled to arbitrarily sized apertures (unlike the errors on the simulations, which are otherwise valid only for point sources).

### 4.3. Catalogs

With this *Paper*, we publish the full SHELA photometric catalog. The catalogs include the IRAC fluxes measured in multiple apertures ( $4''$  and  $6''$  diameter circular apertures, corrected to total magnitudes, isophotal magnitudes, and the “total” (MAG\_AUTO) magnitudes from SExtractor). Errors are estimated from Equation 2 and Table 5 for the linearized number of noise pixels for each object/aperture. In addition, we include a catalog with photometry for the IRAC sources from the SDSS Stripe 82 coadd field (Annis et al. 2014) in *ugriz*, where sources in the SHELA catalog have been matched to the astrometric positions of sources in the SDSS Stripe 82 catalogs using a  $1''$  search radius. Only SDSS sources matched to SHELA sources are included in the catalog, and we include only the closest source in the cases where multiple SDSS sources are located within  $1''$  of a given SHELA source.

The fluxes measured in different apertures will be useful for different studies. In most cases, users will want to use apertures that maximizes the signal-to-noise ratio (SNR) for their objects. For faint, compact sources (including point sources), the maximum SNR occurs for the fluxes measured in the  $4''$  apertures. However, for large, extended sources (much larger than the  $4''$ -diameter aperture), the Isophotal or total aperture may be more appropriate.

Table 6 provides a description of each column name in the tables and the binary FITS tables. Table 7 contains object astrometry and quantities measured from the detection image and weight maps. Tables 8–11 present the SHELA catalogs for the data release. This includes a catalog for the full, combined 3-epoch data (Table 8) and catalogs for each individual epoch (Tables 9–11). The full catalogs are provided as binary tables in Flexible Image Transport System (FITS, Hanisch et al. 2001) format.

The SExtractor flags (FLAGS ( $3.6 \mu\text{m}$ ) and FLAGS ( $4.5 \mu\text{m}$ )) are stored as bits and coded in decimal as the sum of powers of 2 ( $2^{\text{bit}}$ ) for bits that are flagged. Common flag bit values are:

- bit 1** The object has bright neighbors that may bias the photometry, or the object has more than 10% of its pixels flagged as bad or have zero weight;
- bit 2** The object was deblended from another object.

Neither flag bit value is fatal, but users may require a more thorough vetting of these sources depending on their needs. Other, higher (very uncommon in the SHELA catalogs) bit values denote objects whose photometry is dubious. These objects should likely be excluded from use. These bits are available in the SExtractor User’s Manual (v2.13).

**Table 6**  
Column Definitions in SHELA IRAC Catalog

Catalog Column Name <sup>1</sup>	Table Column Name <sup>2</sup>	Description	units	data type
ID	...	Unique ID number from SExtractor for each source in the IRAC catalogs	...	long int
X	...	$x$ -pixel coordinate in IRAC image	pixel	float
Y	...	$y$ -pixel coordinate in IRAC images	pixel	float
RA	RA(J2000)	Right Ascension (J2000) of IRAC source	deg	double
DEC	DEC(J2000)	Declination (J2000) of IRAC source	deg	double
ISOAREA	Isophotal Area	isophotal area of source in detection image	arcsec <sup>2</sup>	float
A	$a$	source semimajor axis	arcsec	float
E	$e$	source ellipticity, $e = 1 - b/a$ , where $b$ is the semiminor axis	...	float
THETA	$\theta$	position angle of the semi-major axis, degrees east from celestial north	deg	float
W3P6	W(3.6)	value of the 3-epoch 3.6 $\mu\text{m}$ weight map at the object's center position <sup>3</sup>	...	float
W4P5	W(4.5)	value of the 3-epoch 4.5 $\mu\text{m}$ weight map at the object's center position <sup>3</sup>	...	float
W3P6_1	W(3.6) <sub>1</sub>	value of the epoch 1, 3.6 $\mu\text{m}$ weight map at the object's center position <sup>3</sup>	...	float
W4P5_1	W(4.5) <sub>1</sub>	value of the epoch 1, 4.5 $\mu\text{m}$ weight map at the object's center position <sup>3</sup>	...	float
W3P6_2	W(3.6) <sub>2</sub>	value of the epoch 2, 3.6 $\mu\text{m}$ weight map at the object's center position <sup>3</sup>	...	float
W4P5_2	W(4.5) <sub>2</sub>	value of the epoch 2, 4.5 $\mu\text{m}$ weight map at the object's center position <sup>3</sup>	...	float
W3P6_3	W(3.6) <sub>3</sub>	value of the epoch 3, 3.6 $\mu\text{m}$ weight map at the object's center position <sup>3</sup>	...	float
W4P5_3	W(4.5) <sub>3</sub>	value of the epoch 3, 4.5 $\mu\text{m}$ weight map at the object's center position <sup>3</sup>	...	float
FLAGS3P6	Flags (3.6 $\mu\text{m}$ )	SExtractor flags for photometry of 3.6 $\mu\text{m}$ image <sup>4</sup>	...	integer
FLAGS4P5	Flags (4.5 $\mu\text{m}$ )	SExtractor flags for photometry of 3.6 $\mu\text{m}$ image <sup>4</sup>	...	integer
F3P6_ISO	$f_{\nu, \text{ISO}}^{(3.6)}$	Isophotal flux density for sources in the 3.6 $\mu\text{m}$ image <sup>4</sup>	$\mu\text{Jy}$	float
F3P6ERR_ISO	$\sigma_{\nu, \text{ISO}}^{(3.6)}$	Error on the 3.6 $\mu\text{m}$ isophotal flux density <sup>4</sup>	$\mu\text{Jy}$	float
F4P5_ISO	$f_{\nu, \text{ISO}}^{(4.5)}$	Isophotal flux density for sources in the 4.5 $\mu\text{m}$ image <sup>4</sup>	$\mu\text{Jy}$	float
F4P5ERR_ISO	$\sigma_{\nu, \text{ISO}}^{(4.5)}$	Error on the 4.5 $\mu\text{m}$ isophotal flux density <sup>4</sup>	$\mu\text{Jy}$	float
F3P6_AUTO	$f_{\nu, \text{AUTO}}^{(3.6)}$	Total flux measured in the Kron aperture for sources in the 3.6 $\mu\text{m}$ image <sup>4</sup>	$\mu\text{Jy}$	float
F3P6ERR_AUTO	$\sigma_{\nu, \text{AUTO}}^{(3.6)}$	Error on the 3.6 $\mu\text{m}$ total flux density <sup>4</sup>	$\mu\text{Jy}$	float
F4P5_AUTO	$f_{\nu, \text{AUTO}}^{(4.5)}$	Total flux measured in a Kron aperture for sources in the 4.5 $\mu\text{m}$ image <sup>4</sup>	$\mu\text{Jy}$	float
F4P5ERR_AUTO	$\sigma_{\nu, \text{AUTO}}^{(4.5)}$	Error on the 4.5 $\mu\text{m}$ total flux density <sup>4</sup>	$\mu\text{Jy}$	float
F3P6_4ARCS	$f_{\nu, 4''}^{(3.6)}$	Flux density measured at 3.6 $\mu\text{m}$ for sources measured in 4''-diameter apertures <sup>4,5</sup>	$\mu\text{Jy}$	float
F3P6ERR_4ARCS	$\sigma_{\nu, 4''}^{(3.6)}$	Error on the flux density at 3.6 $\mu\text{m}$ measured in the 4''-diameter apertures <sup>4</sup>	$\mu\text{Jy}$	float
F4P5_4ARCS	$f_{\nu, 4''}^{(4.5)}$	Flux density measured at 4.5 $\mu\text{m}$ for sources measured in 4''-diameter apertures <sup>4,5</sup>	$\mu\text{Jy}$	float
F4P5ERR_4ARCS	$\sigma_{\nu, 4''}^{(4.5)}$	Error on the flux density at 4.5 $\mu\text{m}$ measured in the 4''-diameter apertures <sup>4</sup>	$\mu\text{Jy}$	float
F3P6_6ARCS	$f_{\nu, 6''}^{(3.6)}$	Flux density measured at 3.6 $\mu\text{m}$ for sources measured in 6''-diameter apertures <sup>4,5</sup>	$\mu\text{Jy}$	float
F3P6ERR_6ARCS	$\sigma_{\nu, 6''}^{(3.6)}$	Error on the flux density at 3.6 $\mu\text{m}$ measured in the 6''-diameter apertures <sup>4</sup>	$\mu\text{Jy}$	float
F4P5_6ARCS	$f_{\nu, 6''}^{(4.5)}$	Flux density measured at 4.5 $\mu\text{m}$ for sources measured in 6''-diameter apertures <sup>4,5</sup>	$\mu\text{Jy}$	float
F4P5ERR_6ARCS	$\sigma_{\nu, 6''}^{(4.5)}$	Error on the flux density at 4.5 $\mu\text{m}$ measured in the 6''-diameter apertures <sup>4</sup>	$\mu\text{Jy}$	float

<sup>1</sup> Column name in binary FITS tables

<sup>2</sup> Column name in Tables 7-11, if different from column name in binary FITS table

<sup>3</sup> The weight map values are proportional to the effective exposure time, with a constant of proportionality  $t_{\text{eff}} = 23.6 \text{ s} \times \text{Weight}$ .

<sup>4</sup> These column names exist in each of the catalogs (combined 3 epoch, and individual epochs) with the same column names.

<sup>5</sup> The flux densities for sources measured in circular apertures have been corrected to total using the aperture corrections in Table 2.

The column descriptions for the binary FITS table and table for the merged SHELA–SDSS Stripe 82 catalog are listed in Table 12. Table 13 presents the photometric data for the merged SHELA–SDSS Stripe 82 catalog.

In the catalogs, objects with no coverage (in a given wavelength and/or epoch) will have weight = 0 and zero flux density and error. These objects also have SExtractor bit=1 set in their flag values. The catalogs and images are available as part of this publication online<sup>11</sup>. It is our intention that these catalogs and images be available through the *Spitzer* Heritage Archive<sup>12</sup>.

<sup>11</sup> <http://drive.google.com/drive/folders/0Bw-jGcrNkzx1dThudFRzM2YrTU0>

<sup>12</sup> We are currently in discussions with *Spitzer* about hosting it on their site, see <http://irsa.ipac.caltech.edu/applications/Spitzer/SHA>, and we expect to migrate all data products to their site while this paper is under review. We provide the data products at the website above to the journal and reviewer as

## 5. A MODICUM OF SCIENCE

### 5.1. Number Counts

Galaxy number counts provide tests of galaxy evolution and cosmology (e.g., Peebles 1993). The number counts are the integral over the luminosity function and distance (redshift), containing the total contribution of galaxies of a given luminosity and distance to the cosmic background emission. The galaxy number counts in the mid-IR are particularly useful as they contain information about stellar-mass growth, dust-obscured populations, and AGN. Galaxy number counts with *Spitzer* have demonstrated the abundance of faint sources attributed to (rest-frame) near-IR and mid-IR emission from distant galaxies and their contribution to the IR background (e.g., Fazio et al. 2004a; Papovich et al. 2004; Dole et al. 2004; Sanders et al. 2007;

a convenience during the review stage.



**Table 12**  
Column Definitions in SHELA/SDSS-matched Catalog

Catalog Column Name <sup>1</sup>	Table Column Name <sup>2</sup>	Description	units	data type
SHELA_ID	ID	Unique ID from the SHELA catalog in Table 7	...	long int
SDSS_ID	SDSS ID	ID of object in the SDSS catalogs	...	64-bit long int
SDSS_RA	SDSS RA	Right ascension (J2000) of object in the SDSS catalog	deg	double
SDSS_DEC	SDSS DEC	Declination (J2000) of object in the SDSS catalog	deg	double
TYPE	...	Object type from SDSS catalog <sup>3</sup>	...	int
FLAGS	SDSS FLAGS	SDSS Flags for the object	...	64-bit long int
U	$u$	SDSS $u$ total AB magnitude corrected for Galactic extinction	mag	float
UERR	$\sigma_u$	uncertainty on SDSS $g$ AB magnitude	mag	float
G	$g$	SDSS $g$ total AB magnitude corrected for Galactic extinction	mag	float
GERR	$\sigma_g$	uncertainty on SDSS $g$ AB magnitude	mag	float
R	$r$	SDSS $r$ total AB magnitude corrected for Galactic extinction	mag	float
RERR	$\sigma_r$	uncertainty on SDSS $r$ AB magnitude	mag	float
I	$i$	SDSS $i$ total AB magnitude corrected for Galactic extinction	mag	float
IERR	$\sigma_i$	uncertainty on SDSS $i$ AB magnitude	mag	float
Z	$z$	SDSS $z$ total AB magnitude corrected for Galactic extinction	mag	float
ZERR	$\sigma_z$	uncertainty on SDSS $z$ AB magnitude	mag	float

<sup>1</sup> Column name in binary FITS tables

<sup>2</sup> Column name in Table 13, if different from column name in binary FITS table

<sup>3</sup> The most common type values are TYPE=3 for Galaxy or TYPE=6 for Star.

Ashby et al. 2009, 2013b; Mauduit et al. 2012).

Figure 19 shows the IRAC 3.6 and 4.5  $\mu\text{m}$  number counts for the full SHELA data. The raw counts (uncorrected for completeness, see § 4.1) are provided in Table 14. The SHELA counts agree well with previous measurements (e.g., Fazio et al. 2004b; Sanders et al. 2007). At bright magnitudes ( $AB \lesssim 18$ ) the counts follow roughly the expected contribution from Galactic stars (e.g., Fazio et al. 2004b; Ashby et al. 2013b, and references therein). The SHELA counts show a slight excess of bright counts compared to the data in Fazio et al. and Sanders et al. This is likely a result of the different Galactic sightlines compared to the fields of those studies, and therefore the excess number density of Galactic stars is not surprising.

The counts provide an independent measure of the completeness of the SHELA IRAC catalogs. The SHELA counts in Figure 19 include the completeness corrections derived in § 4.1. These counts agree with those derived from deeper IRAC images measured by Fazio et al. (2004b) and Sanders et al. (2007), as illustrated in Figure 19, at least down to  $m_{AB} \sim 22.5$  mag where the SHELA data are more than 50% complete. At fainter magnitudes, the completeness corrections for the SHELA data are significantly higher and the uncertainties on the completeness corrections dominate the counts (and Eddington-type biases are most severe). Therefore, we have confidence in the completeness corrections and the number counts down to the 50% completeness limit.

### 5.2. Time-Variable Objects

The combination of the large area and multi-epoch nature of the SHELA dataset allows for the identification of sources whose brightness varies across the  $\sim 6$  and 12 month baselines in time. This includes rare sources that show large changes in brightness, and sources with high proper motion (see also, Ashby et al. 2009).

As an illustration, we selected objects from the SHELA catalog that are detected in both the combined 3.6 and 4.5  $\mu\text{m}$  data, have coverage in all three observing epochs, but vary by more than 2.5 mag (a factor of 10 in brightness) between any two observing epochs. There are 291 objects in the SHELA field that satisfy these requirements with  $[3.6] \leq 20.5$  AB mag or  $[4.5] \leq 20.5$  AB mag. An inspection of these objects


shows they are all consistent with point sources, with several “double” (resolved, or multiple component) objects and some objects that appear to show astrometric centroid shifts between the 3.6 and 4.5  $\mu\text{m}$  image (which would imply very high proper motions, indicative of asteroids). Figure 20 shows four objects that appear in only a single observing epoch. Because such objects make it into the final, three-epoch, combined catalog, care should be made in selecting object samples that require objects with no (significant) temporal variation in brightness.

### 5.3. The Relation between the Scale Radius and Mass of Dark-Matter Halos


The ACT survey includes the SHELA IRAC footprint, and its catalog includes SZ emission from distant ( $z < 1$ ) clusters in the Stripe 82 field (Hasselfield et al. 2013). The thermal SZ effect is a decrement in the emission from the CMB owing to the presence of a massive (virialized) galaxy cluster along the line of sight. The hot ( $T \sim 10^7 - 10^8$  K) ICM gas associated with the galaxy cluster causes inverse Compton scattering of the CMB photons, leaving a distortion in the direction of the cluster. The strength of the distortion is proportional to the line-of-sight integral of the thermal pressure (the Compton  $y$  parameter), which correlates with the total mass ( $M_{500}$ ) associated with the galaxy cluster (e.g., Vikhlinin et al. 2009; Marrone et al. 2012; Sifón et al. 2013).

Five of the ACT clusters from Hasselfield et al. (2013) fall in the SHELA/IRAC footprint (ACT CLJ0059.1-0049, ACT CLJ0127.2+0020, ACT CLJ0119.9+0055, ACT CLJ0058.0+0030, ACT CLJ0104.8+0002). The IRAC data probe the amount of starlight associated with the galaxies in these clusters, and measure the galaxy spatial distribution. The combination of IRAC and ACT data therefore allows us to study the structural size of the dark matter halo (as traced by the galaxies in the IRAC image) and compare it with the halo mass as estimated from the SZ signal.

Studies have shown that the surface density of satellites roughly trace the distribution of dark matter (e.g., Tal et al. 2012; Kawinwanichakij et al. 2014; van der Burg et al. 2015) predicted by the density profile of the dark matter from numerical simulations (e.g., Navarro et al. 1996, NFW hereafter). The Spitzer/IRAC data allow the measurement of




**Figure 19.** Differential number counts of IRAC sources in the SHELDA field in bins of 0.2 mag. The left panel shows the results for the IRAC 3.6  $\mu\text{m}$  data. The right panel shows the results for the 4.5  $\mu\text{m}$  data. In each panel the heavy black histogram shows the IRAC counts with no correction for completeness. The error bars show Poisson uncertainties on the number counts. The gray line shows the counts corrected for incompleteness. For comparison the dashed line shows counts from S-COSMOS (Sanders et al. 2007) and the dot-dashed line shows counts from the IRAC GTO data (Fazio et al. 2004a).



**Figure 20.** Examples of sources that vary in brightness by more than 2.5 magnitudes between the different SHELDA observing epochs. Each set of panels shows the 3.6 and 4.5  $\mu\text{m}$  images for epochs 1, 2, and 3 for 4 sources in the SHELDA catalog. The object catalog IDs are given as the title for each set of plots.






**Figure 21.** The left panel shows the SHEL A IRAC  $3.6 \mu\text{m}$  image of the ACT SZ-selected cluster CLJ0058.0+0030 at  $z = 0.76$  (Hasselfield et al. 2013). The right panel shows the projected surface-density distribution of galaxies centered on the peak of the SZ signal of the cluster. The surface density is the number of galaxies in the SHEL A IRAC data measured in concentric annuli centered on the cluster, corrected for the average density of galaxies in random apertures in the IRAC image. The surface density is consistent with a projected NFW profile with scale radius  $0.51 \pm 0.14$  arcmin.

**Table 14**  
SHELA IRAC Number Counts

$m_{AB}$ (mag) (1)	IRAC 3.6 $\mu\text{m}$		IRAC 4.5 $\mu\text{m}$	
	$dN/dm$ ( $\text{mag}^{-1} \text{deg}^{-2}$ ) (2)	Error ( $\text{mag}^{-1} \text{deg}^{-2}$ ) (3)	$dN/dm$ ( $\text{mag}^{-1} \text{deg}^{-2}$ ) (4)	Error ( $\text{mag}^{-1} \text{deg}^{-2}$ ) (5)
12.0	7.3	1.1	3.8	0.8
12.2	7.0	1.0	3.5	0.7
12.4	9.9	1.2	6.7	1.0
12.6	13.8	1.4	7.8	1.1
12.8	20.8	1.8	14.1	1.5
13.0	24.3	1.9	22.6	1.9
13.2	31.5	2.2	19.3	1.7
13.4	36.9	2.4	21.4	1.8
13.6	35.4	2.3	22.9	1.9
13.8	44.4	2.6	31.0	2.2
14.0	52.0	2.8	34.2	2.3
14.2	61.2	3.1	40.7	2.5
14.4	67.2	3.2	47.7	2.7
14.6	78.6	3.5	56.4	2.9
14.8	93.5	3.8	62.0	3.1
15.0	106.2	4.0	77.2	3.4
15.2	117.2	4.2	84.7	3.6
15.4	140.0	4.6	95.8	3.8
15.6	168.2	5.1	116.6	4.2
15.8	175.8	5.2	134.7	4.5
16.0	215.1	5.7	152.7	4.8
16.2	243.3	6.1	186.2	5.3
16.4	287.1	6.6	219.3	5.8
16.6	335.4	7.1	250.2	6.2
16.8	394.6	7.7	306.9	6.8
17.0	469.0	8.4	358.1	7.4
17.2	561.0	9.2	429.1	8.1
17.4	693.5	10.0	533.6	9.0
17.6	853.4	11.0	643.2	9.9
17.8	1096.0	13.0	809.5	11.0
18.0	1417.0	15.0	1032.0	13.0
18.2	1839.0	17.0	1324.0	14.0
18.4	2427.0	19.0	1697.0	16.0
18.6	3138.0	22.0	2260.0	19.0
18.8	4005.0	25.0	2952.0	21.0
19.0	5088.0	28.0	3881.0	24.0
19.2	6322.0	31.0	5012.0	28.0
19.4	7523.0	34.0	6372.0	31.0
19.6	8904.0	37.0	7931.0	35.0
19.8	10390.0	40.0	9481.0	38.0
20.0	11780.0	42.0	11180.0	41.0
20.2	13040.0	45.0	12680.0	44.0
20.4	14240.0	47.0	14110.0	46.0
20.6	15330.0	48.0	15310.0	48.0
20.8	16760.0	50.0	16620.0	50.0
21.0	18400.0	53.0	18080.0	52.0
21.2	19810.0	55.0	19630.0	55.0
21.4	20950.0	56.0	20900.0	56.0
21.6	21420.0	57.0	21210.0	57.0
21.8	21360.0	57.0	21100.0	57.0
22.0	20740.0	56.0	20730.0	56.0
22.2	19680.0	55.0	19610.0	55.0
22.4	17840.0	52.0	18120.0	52.0
22.6	15140.0	48.0	15670.0	49.0
22.8	11690.0	42.0	12490.0	44.0
23.0	8166.0	35.0	9101.0	37.0

**Note.** — (1) magnitude of number count bin, (2) number counts at 3.6  $\mu\text{m}$ , (3) Poisson error on 3.6  $\mu\text{m}$  number counts, (4) number counts at 4.5  $\mu\text{m}$ , (5) Poisson error on 4.5  $\mu\text{m}$  number counts. Note that the counts are not corrected for completeness. To do so requires dividing by the magnitude-dependent completeness corrections in Table 4.

the radial distribution of galaxies, and therefore a tracer of the dark matter density distribution. We use the method from Kawinwanichakij et al. (2014). We count the number of galaxies in the SHELA IRAC catalog with  $17 < m_{3.6} < 22$  mag in concentric annuli centered on each cluster. To correct for the background, we measure the average (median)



**Figure 22.** The evolution of the ratio between the NFW-profile scale radius,  $r_s$  and halo mass,  $M_{200}$ . The boxes and error bars show values derived for the ACT-selected clusters in SHELA. The scale radii are measured by fitting projected NFW profiles to the surface density of galaxies in each ACT cluster. The asterisks show the predicted ratio of  $r_s/M_{200}^{0.45}$  for galaxy halos from NFW96. The hatched swaths show the expected ratio for halos of  $\log M_{200}/M_\odot = 13, 14,$  and  $15$  (as labeled in the figure legend) including the scatter in halo concentration using the relations in Diemer & Kravtsov (2015). The data are consistent with a near unevolving ratio of  $r_s/M_{200}^{0.45}$  over a large baseline in redshift, with values consistent with DM halo scaling relations as expected from the distribution of predicted ratios for a CDM-type cosmology.

number of galaxies in each annulus for  $10^4$  randomly placed apertures around the SHELA image (taking care to avoid the image edges). We then measure the radial profile, and fit the projected NFW profile (Bartelmann 1996) using two parameters, the NFW scale angular radius,  $\theta_s$ , and a normalization. The results from these fits for the NFW scale radii for each cluster are given in Table 15.

Figure 21 shows the distribution of galaxies centered on one of the ACT clusters in SHELA, ACT CLJ0058.0+0030 at  $z = 0.76$ . The galaxy distribution has been corrected statistically for the galaxies associated with the field as discussed above. The figure shows that the surface density of galaxies follows a projected NFW profile with a best fit scale radius,  $\theta_s = 0.51 \pm 0.14$  arcmin, which corresponds to a physical scale radius of  $r_s = 223 \pm 65$  kpc (for  $h = 0.7$ ,  $\Omega_m = 0.3$ , and  $\Omega_\Lambda = 0.7$ ).

The combination of data on the radial distribution of cluster galaxies and measures of the total mass of the clusters is a potentially powerful way to study properties of dark matter halos. For an NFW profile, the scale radius is expected to increase with halo mass as  $r_s \sim M_{200}^{0.45}$  (Navarro et al. 1996), where  $M_{200}$  is the mass within a radius where the density is 200 times the critical density. Simulations predict that this relation is expected to remain roughly constant with redshift (e.g., Bullock et al. 2001; Eke et al. 2001; Diemer & Kravtsov 2015).

Figure 22 shows the ratio of  $r_s/M_{200}^{0.45}$  for the five ACT SZ clusters in the SHELA field as a function of their redshift, where the  $r_s$  values come from the projected NFW profile fits to the radial distributions of galaxies for each cluster from the IRAC catalogs (as described for CLJ0058.0+0030 in the previous paragraph) and the  $M_{200}$  values come from measurements of the SZ  $y$ -parameter (Hasselfield et al. 2013, where we have adjusted the  $M_{500}$  values by 0.1 dex to convert them to



**Table 15**  
Measures of Scale Radii of NFW profiles in ACT SZ Clusters in SHELA

Cluster	$z$	$M_{500}$ ( $10^{14} M_{\odot}$ )	$\theta_s$ (arcmin)	$r_s$ (kpc)
(1)	(2)	(3)	(4)	(5)
ACT CLJ0104.8+0002	0.28	$2.6 \pm 0.9$	$1.1 \pm 0.4$	$280 \pm 100$
ACT CLJ0127.2+0020	0.37	$3.3 \pm 0.9$	$0.67 \pm 0.24$	$205 \pm 72$
ACT CLJ0119.9+0055	0.72	$3.3 \pm 0.8$	$0.72 \pm 0.22$	$311 \pm 96$
ACT CLJ0058.0+0030	0.76	$3.2 \pm 0.8$	$0.51 \pm 0.15$	$223 \pm 65$
ACT CLJ0059.1-0049	0.77	$5.2 \pm 0.9$	$0.72 \pm 0.19$	$318 \pm 83$

**Note.** — (1) Cluster designation (from Hasselfield et al. 2013), (2) estimated redshift (from Hasselfield et al. 2013), (3) total cluster mass derived from the SZ signal (from Hasselfield et al. 2013), (4) angular scale radius  $\theta_s$  of the projected NFW profile fit to the background-corrected surface density of galaxies in the SHELA IRAC data centered on each cluster, (5) scale radius converted to physical units assuming  $h = 0.7$ ,  $\Omega_m = 0.3$ , and  $\Omega_{\Lambda} = 0.7$ .

$M_{200}$ ). Figure 22 also shows the values of  $r_s/M_{200}^{0.45}$  for the simulated model halos from Navarro et al. (1996), for halos ranging from  $M_{200} = 10^{13} - 10^{15} M_{\odot}$ . The values range from  $r_s/M_{200}^{0.45} \simeq 0.1 - 0.2$  Mpc /  $(10^{14} M_{\odot})^{0.45}$ , and these agree remarkably well with the observations for the 5 ACT-SZ clusters. These are consistent with the expected evolution of more modern simulations for a halo of mass  $10^{13} - 10^{15} M_{\odot}$  (Diemer & Kravtsov 2015). The observations show indications that the shape of the dark-matter profiles has only a weak dependence on mass and redshift in accordance with predictions from  $\Lambda$ CDM (Bullock et al. 2001).

## 6. SUMMARY

We presented the *Spitzer* IRAC imaging at 3.6 and 4.5  $\mu$ m of the SHELA survey, a *Spitzer* Exploratory program which covers a  $\approx 24$  deg<sup>2</sup> field within the footprint of HETDEX. This field has a rich set of multiwavelength imaging, including the SDSS Stripe 82 imaging, DES, CTIO/DECam, NEWFIRM *K*-band, *Herschel* far-IR imaging, and Chandra and *XMM-Newton* X-ray imaging.

The HETDEX survey will obtain redshifts in this field for  $\sim 200,000$  galaxies at  $1.9 < z < 3.5$  based on Lyman- $\alpha$  emission (covering a volume of 0.5 Gpc<sup>3</sup>), and redshifts for an additional  $\sim 200,000$  galaxies at  $z < 0.5$  based on their [O II] emission. The SHELA IRAC data are sensitive to galaxies with stellar masses down to  $\simeq 2 \times 10^{10} M_{\odot}$  through the redshift range of Lyman- $\alpha$  probed by HETDEX. Thus, the combination of the HETDEX spectroscopy data, ground-based optical/near-IR imaging, and the SHELA IRAC data allow the study of the relationship between structure formation, galaxy stellar mass, dark halo mass, and environment during over a large range of cosmic history.

In this *Paper* we discussed the properties of the SHELA IRAC data, including the data acquisition, reduction, validation, and source catalogs. The imaging includes three observing epochs separated by approximately 6 months between epochs. The combined three-epoch dataset covers 24.2 deg<sup>2</sup> with an exposure time of at least  $\approx 200$  s. Our tests show the images and catalogs are 80% (50%) complete to limiting magnitudes of 22.0 (22.6) AB mag in the detection image, constructed from the weighted-sum of the IRAC 3.6 and 4.5  $\mu$ m images. The catalogs reach limiting ( $1\sigma$ ) sensitivities of 1.1  $\mu$ Jy in each IRAC channel. The photometric accuracy is consistent with AllWISE with essentially no different between the [3.6] and W1 bands and a possible 0.02 mag offset between [4.5] and W2 bands.

The astrometric solution of SHELA is tied to SDSS DR7, where the astrometric uncertainty is  $< 0''.2$ , comparable to the uncertainty in the SDSS catalogs. The astrometric solutions

are accurate compared to 2MASS, but show a (possibly) non-negligible offset compared to the AllWISE approaching one-tenth of an arcsecond.

The IRAC data enable a broad range of scientific explorations, including studies of galaxy and AGN evolution, and the formation of large-scale structure. As a demonstration of science, we present IRAC number counts, examples of highly temporally variable sources, and galaxy surface density profiles of rich galaxy clusters. At faint magnitudes, the source number counts are consistent with other IRAC datasets, which provides confidence in our estimated completeness corrections. At bright magnitudes we observe a possible excess of counts, which we attribute to variations in the surface density of Galactic stars.

We use a sample of five ACT SZ-selected galaxy clusters between  $0.2 < z < 0.8$  to study the relation between cluster mass (traced by the SZ Compton parameter) and the scale radius,  $r_s$ , of the cluster halos as traced by the surface distribution of galaxies as measured from the SHELA IRAC data. All clusters show galaxy surface densities in agreement with a projected NFW halo, with a ratio of  $r_s/M_{200}^{0.45}$  that is consistent with simulations of dark matter halos and is unevolving in redshift, as predicted by  $\Lambda$ CDM models.

In the spirit of *Spitzer* Exploratory programs we provide all images and catalogs as part of this *Paper* (see footnote 11). We describe the source IRAC catalogs and imaging products released for the SHELA data, which will be available through IRSA.

We acknowledge useful conversations, comments and suggestions, especially from Matt Ashby, Ivo Labbé, Gordon Richards, Nick Ross, Eli Rykoff, Louis Strigari, and John Timlin. CP thanks the Space Telescope Science Institute for its hospitality and for providing a scientifically conducive atmosphere during the completion of this work. This work is based on observations made with the *Spitzer* Space Telescope, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. This work is supported by the National Science Foundation through grant AST-1413317. We acknowledge generous support from the Texas A&M University and the George P. and Cynthia Woods Institute for Fundamental Physics and Astronomy. RC, CG, and GZ acknowledge support from the Institute for Gravitation and the Cosmos. The Institute for Gravitation and the Cosmos is supported by the Eberly College of Science and the Office of the Senior Vice President for Research at the Pennsylvania State University.

This work made use of data from SDSS. Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan

Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is <http://www.sdss.org>.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration.

## REFERENCES

- Abazajian, K. N., Adelman-McCarthy, J. K., Agüeros, M. A., et al. 2009, *ApJS*, 182, 543
- Adams, J. J., Blanc, G. A., Hill, G. J., et al. 2011, *ApJS*, 192, 5
- Alam, S., Albareti, F. D., Allende Prieto, C., et al. 2015, *ApJS*, 219, 12
- Annis, J., Soares-Santos, M., Strauss, M. A., et al. 2014, *ApJ*, 794, 120
- Ashby, M. L. N., Stern, D., Brodwin, M., et al. 2009, *ApJ*, 701, 428
- Ashby, M. L. N., Willner, S. P., Fazio, G. G., et al. 2013a, *ApJ*, 769, 80
- Ashby, M. L. N., Stanford, S. A., Brodwin, M., et al. 2013b, *ApJS*, 209, 22
- Ashby, M. L. N., Willner, S. P., Fazio, G. G., et al. 2015, *ApJS*, 218, 33
- Bartelmann, M. 1996, *A&A*, 313, 697
- Beers, T. C., Flynn, K., & Gebhardt, K. 1990, *AJ*, 100, 32
- Behroozi, P. S., Conroy, C., & Wechsler, R. H. 2010, *ApJ*, 717, 379
- Behroozi, P. S., Wechsler, R. H., & Conroy, C. 2013, *ApJ*, 762, L31
- Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
- Bertin, E., Mellier, Y., Radovich, M., et al. 2002, in *Astronomical Society of the Pacific Conference Series*, Vol. 281, *Astronomical Data Analysis Software and Systems XI*, ed. D. A. Bohlender, D. Durand, & T. H. Handley, 228
- Blanc, G. A., Lira, P., Barrientos, L. F., et al. 2008, *ApJ*, 681, 1099
- Blanc, G. A., Adams, J. J., Gebhardt, K., et al. 2011, *ApJ*, 736, 31
- Bullock, J. S., Kolatt, T. S., Sigad, Y., et al. 2001, *MNRAS*, 321, 559
- Cutri, R. M., Wright, E. L., Conrow, T., et al. 2013, *Explanatory Supplement to the AllWISE Data Release Products*, Tech. rep.
- Diemer, B., & Kravtsov, A. V. 2015, *ApJ*, 799, 108
- Dole, H., Le Floch, E., Pérez-González, P. G., et al. 2004, *ApJS*, 154, 87
- Eisenhardt, P. R. M., Brodwin, M., Gonzalez, A. H., et al. 2008, *ApJ*, 684, 905
- Eke, V. R., Navarro, J. F., & Steinmetz, M. 2001, *ApJ*, 554, 114
- Fazio, G. G., Ashby, M. L. N., Barmby, P., et al. 2004a, *ApJS*, 154, 39
- Fazio, G. G., Hora, J. L., Allen, L. E., et al. 2004b, *ApJS*, 154, 10
- Finkelstein, S. L., Hill, G. J., Gebhardt, K., et al. 2011, *ApJ*, 729, 140
- Gawiser, E., van Dokkum, P. G., Herrera, D., et al. 2006, *ApJS*, 162, 1
- Hagen, A., Ciardullo, R., Gronwall, C., et al. 2014, *ApJ*, 786, 59
- Hanisch, R. J., Farris, A., Greisen, E. W., et al. 2001, *A&A*, 376, 359
- Hasselfield, M., Hilton, M., Marriage, T. A., et al. 2013, *JCAP*, 7, 8
- Hill, G. J., Gebhardt, K., Komatsu, E., et al. 2008, in *Astronomical Society of the Pacific Conference Series*, Vol. 399, *Panoramic Views of Galaxy Formation and Evolution*, ed. T. Kodama, T. Yamada, & K. Aoki, 115
- Ivezić, Ž., Smith, J. A., Miknaitis, G., et al. 2007, *AJ*, 134, 973
- Jarrett, T. H., Cohen, M., Masci, F., et al. 2011, *ApJ*, 735, 112
- Kawinwanichakij, L., Papovich, C., Quadri, R. F., et al. 2014, *ApJ*, 792, 103
- Komatsu, E., Smith, K. M., Dunkley, J., et al. 2011, *ApJS*, 192, 18
- Kravtsov, A., Vikhlinin, A., & Meshcheryakov, A. 2014, *ApJ*, submitted, arXiv:1401.7329
- Kurucz, R. L. 1993, *SYNTH* spectrum synthesis programs and line data
- Labbé, I., Franx, M., Rudnick, G., et al. 2003, *AJ*, 125, 1107
- Labbé, I., Oesch, P. A., Bouwens, R. J., et al. 2013, *ApJ*, 777, L19
- LaMassa, S. M., Urry, C. M., Glikman, E., et al. 2013a, *MNRAS*, 432, 1351
- LaMassa, S. M., Urry, C. M., Cappelluti, N., et al. 2013b, *MNRAS*, 436, 3581
- Lonsdale, C. J., Smith, H. E., Rowan-Robinson, M., et al. 2003, *PASP*, 115, 897
- Marrone, D. P., Smith, G. P., Okabe, N., et al. 2012, *ApJ*, 754, 119
- Matsuura, M., Zijlstra, A. A., van Loon, J. T., et al. 2005, *A&A*, 434, 691
- Matsuura, M., Bernard-Salas, J., Lloyd Evans, T., et al. 2014, *MNRAS*, 439, 1472
- Mauduit, J.-C., Lacy, M., Farrah, D., et al. 2012, *PASP*, 124, 714
- Moster, B. P., Naab, T., & White, S. D. M. 2013, *MNRAS*, 428, 3121
- Moster, B. P., Somerville, R. S., Maulbetsch, C., et al. 2010, *ApJ*, 710, 903
- Muzzin, A., Wilson, G., Demarco, R., et al. 2013, *ApJ*, 767, 39
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1996, *ApJ*, 462, 563
- Oke, J. B., & Gunn, J. E. 1983, *ApJ*, 266, 713
- Papovich, C. 2008, *ApJ*, 676, 206
- Papovich, C., Dole, H., Egami, E., et al. 2004, *ApJS*, 154, 70
- Papovich, C., Labbé, I., Quadri, R., et al. 2015, *ApJ*, 803, 26
- Peebles, P. J. E. 1993, *Principles of physical cosmology* (Princeton, NJ: Princeton University Press)
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2014, *A&A*, 571, A16
- Quadri, R., van Dokkum, P., Gawiser, E., et al. 2007, *ApJ*, 654, 138
- Reach, W. T., Megeath, S. T., Cohen, M., et al. 2005, *PASP*, 117, 978
- Richards, G. T., Myers, A. D., Peters, C. M., et al. 2015, *ApJS*, 219, 39
- Sanders, D. B., Salvato, M., Aussel, H., et al. 2007, *ApJS*, 172, 86
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Sifón, C., Menanteau, F., Hasselfield, M., et al. 2013, *ApJ*, 772, 25
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Somerville, R. S., & Davé, R. 2015, *ARA&A*, 53, 51
- Springel, V., White, S. D. M., Jenkins, A., et al. 2005, *Nature*, 435, 629
- Stetson, P. B. 1987, *PASP*, 99, 191
- Tal, T., Wake, D. A., & van Dokkum, P. G. 2012, *ApJ*, 751, L5
- Timlin, J. D., Ross, N. P., Richards, G. T., et al. 2015, in preparation
- Tinker, J., Kravtsov, A. V., Klypin, A., et al. 2008, *ApJ*, 688, 709
- van der Burg, R. F. J., Hoekstra, H., Muzzin, A., et al. 2015, *A&A*, 577, A19
- Viero, M. P., Asboth, V., Roseboom, I. G., et al. 2014, *ApJS*, 210, 22
- Vikhlinin, A., Burenin, R. A., Ebeling, H., et al. 2009, *ApJ*, 692, 1033
- Weinberg, D. H., Davé, R., Katz, N., & Hernquist, L. 2004, *ApJ*, 601, 1
- Werner, M. W., Roellig, T. L., Low, F. J., et al. 2004, *ApJS*, 154, 1
- Whitaker, K. E., Labbé, I., van Dokkum, P. G., et al. 2011, *ApJ*, 735, 86
- Wilson, G., Muzzin, A., Yee, H. K. C., et al. 2009, *ApJ*, 698, 1943

**Table 7**  
Preamble for all SHELA IRAC Catalogs

ID	X	Y	RA(J2000)	DEC(J2000)	Isophotal Area	$a$	$e$	$\theta$	W(3.6)	W(4.5)	W(3.6) <sub>1</sub>	W(4.5) <sub>1</sub>	W(3.6) <sub>2</sub>	W(4.5) <sub>2</sub>	W(3.6) <sub>3</sub>	W(4.5) <sub>3</sub>
(1)	(pixel) (2)	(pixel) (3)	(deg) (4)	(deg) (5)	(arcsec <sup>2</sup> ) (6)	(arcsec) (7)	(8)	(deg) (9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)
100020	45386.8	1082.6	17.701277	-1.132968	125.4	1.9	0.04	35.3	2.92	6.03	0.00	1.02	2.92	3.00	0.00	2.00
100021	18859.7	1115.1	23.590947	-1.125582	7.0	0.8	0.25	-13.1	8.01	10.76	3.00	3.01	2.02	4.77	2.99	2.99
100022	19340.1	1112.4	23.484482	-1.126288	12.2	1.1	0.26	-0.7	6.32	6.03	3.28	2.99	0.00	0.00	3.04	3.05
100023	19528.0	1112.8	23.442827	-1.126232	12.2	1.1	0.46	-29.2	6.03	7.06	3.04	3.03	0.00	0.00	2.99	4.03
100024	56605.8	1102.2	15.221331	-1.125113	26.9	1.2	0.11	-53.5	4.08	9.01	0.00	2.98	4.08	3.03	0.00	3.01
100025	22941.5	1112.9	22.685727	-1.126878	7.0	0.8	0.31	-41.1	11.12	9.05	5.03	2.99	3.04	3.01	3.05	3.05
100026	21391.8	1115.2	23.029557	-1.126084	3.8	0.6	0.25	-47.4	3.04	6.79	0.00	2.77	3.04	0.00	0.00	4.01
100027	45127.6	1104.6	17.758739	-1.128140	25.6	1.3	0.14	-32.0	7.18	7.15	3.00	4.11	4.17	3.03	0.00	0.00
100028	28402.5	1118.5	21.473082	-1.126254	5.8	0.8	0.56	-18.8	2.99	6.15	0.00	3.14	2.99	0.00	0.00	3.01
100029	35643.7	1111.3	19.864062	-1.127908	12.2	1.0	0.19	-32.0	0.00	8.06	0.00	5.05	0.00	0.00	0.00	3.01
100030	22402.3	1098.1	22.805387	-1.130069	30.1	1.9	0.42	-65.9	8.00	9.09	3.02	3.11	2.99	2.99	1.99	2.99
100031	26946.7	1113.5	21.796466	-1.127243	6.4	0.8	0.30	-89.2	3.00	3.01	3.00	3.01	0.00	0.00	0.00	0.00
100032	8436.4	1095.7	25.894587	-1.126576	62.7	2.1	0.23	-7.7	9.07	6.03	6.03	2.98	0.00	0.00	3.04	3.05
100033	28542.3	1118.6	21.442033	-1.126240	3.2	0.6	0.39	-16.6	2.99	6.02	0.00	3.03	2.99	0.00	0.00	2.99
100034	27031.2	1113.5	21.777696	-1.127268	5.1	0.9	0.55	67.7	3.00	3.38	3.00	3.38	0.00	0.00	0.00	0.00
100035	29155.7	1118.3	21.305746	-1.126346	1.9	0.5	0.40	44.8	14.88	6.01	3.02	3.00	8.84	0.00	3.02	3.00
100036	41240.4	1091.2	18.621071	-1.131813	84.5	1.8	0.24	34.9	9.07	10.76	3.03	2.41	3.04	3.02	3.00	5.34
100037	33861.8	1108.3	20.259993	-1.128661	16.6	1.1	0.16	-23.1	9.04	8.14	3.03	2.11	2.99	2.99	3.02	3.04
100038	40805.0	1105.7	18.717715	-1.128666	21.8	1.1	0.14	59.5	7.94	4.25	3.03	2.03	2.88	0.00	2.02	2.22
100039	50844.3	1103.6	16.492991	-1.126869	21.1	1.5	0.43	-82.5	5.17	6.04	2.15	3.00	0.00	0.00	3.02	3.04
100040	34697.3	1108.7	20.074340	-1.128531	14.7	1.1	0.17	79.3	0.00	4.85	0.00	3.01	0.00	0.00	0.00	1.85
100041	35355.1	1108.5	19.928174	-1.128544	19.2	1.1	0.14	-62.1	0.00	6.08	0.00	3.06	0.00	0.00	0.00	3.02

**Note.** — This is a portion of the full table to provide form and guidance. 1. Unique object ID number, 2. central X pixel coordinate, 3. central Y pixel coordinate, 4. object right ascension (J2000) in decimal degrees, 5. object declination (J2000) in decimal degrees, 6. Isophotal area in the detection (combined 2 band, 3 epoch) image, 7. semimajor axis in the detection image, 8. ellipticity measured in the detection image, defined as  $e = 1 - b/a$ , where  $b$  and  $a$  are the semiminor and semimajor axes, respectively, 9. position angle measured in the detection image (degrees E from N), 10–17. values of the weight maps in the images at the location of the object. The weight map is proportional to the exposure time map. 10–11. values in combined 3.6 and 4.5  $\mu\text{m}$  image weight maps, respectively. 12.–17. values in the 3.6 and 4.5  $\mu\text{m}$  weight maps for the individual epochs, respectively.



**Table 8**  
Photometry for Combined, 3 Epoch SHELA IRAC Catalogs

ID	Flags (3.6 $\mu$ m)	Flags (4.5 $\mu$ m)	$f_{\nu,ISO}^{(3.6)}$ ( $\mu$ Jy)	$\sigma_{ISO}^{(3.6)}$ ( $\mu$ Jy)	$f_{\nu,AUTO}^{(3.6)}$ ( $\mu$ Jy)	$\sigma_{AUTO}^{(3.6)}$ ( $\mu$ Jy)	$f_{\nu,4''}^{(3.6)}$ ( $\mu$ Jy)	$\sigma_{4''}^{(3.6)}$ ( $\mu$ Jy)	$f_{\nu,6''}^{(3.6)}$ ( $\mu$ Jy)	$\sigma_{6''}^{(3.6)}$ ( $\mu$ Jy)	$f_{\nu,ISO}^{(4.5)}$ ( $\mu$ Jy)	$\sigma_{ISO}^{(4.5)}$ ( $\mu$ Jy)	$f_{\nu,AUTO}^{(4.5)}$ ( $\mu$ Jy)	$\sigma_{AUTO}^{(4.5)}$ ( $\mu$ Jy)	$f_{\nu,4''}^{(4.5)}$ ( $\mu$ Jy)	$\sigma_{4''}^{(4.5)}$ ( $\mu$ Jy)	$f_{\nu,6''}^{(4.5)}$ ( $\mu$ Jy)	$\sigma_{6''}^{(4.5)}$ ( $\mu$ Jy)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)
100020	2	2	454	11.9	452	12	334	9.53	408	10.6	525	12.7	525	12.8	392	10.3	470	11.3
100021	0	0	2.86	1.02	10.2	3.18	4.72	1.36	5.76	1.83	2.98	1.03	3.81	2.9	5.06	1.39	4.37	1.72
100022	2	2	5.7	1.44	6.89	3.22	7.33	1.6	8.93	2.05	5.09	1.38	8.42	3.29	6.96	1.56	7.03	1.92
100023	1	1	3.72	1.24	2.87	1.46	2.93	1.17	2.42	1.56	5.71	1.44	7.46	1.84	6.78	1.55	7.35	1.94
100024	0	0	43.5	3.66	45	4.02	48.5	3.7	49.3	3.89	26.7	2.98	26.8	3.35	31.5	3.01	30.8	3.18
100025	0	0	3.24	1.06	6.68	2.67	5.23	1.41	6.22	1.86	2.92	1.02	4.03	2.53	4.25	1.31	3.16	1.63
100026	0	0	2.05	0.817	3.47	1.17	4.99	1.38	4.78	1.75	2.47	0.883	3.54	1.18	4.7	1.36	3.96	1.69
100027	0	0	19	2.58	20	3.17	19.7	2.43	20.8	2.72	23.7	2.82	25.5	3.4	26.6	2.78	27	3.01
100028	0	0	1.68	0.806	1.9	1.32	3.78	1.26	3.73	1.67	3.36	1.05	4.2	1.54	4.68	1.35	4.35	1.72
100029	1	0	0	0	0	0	0	0	0	0	9.3	1.75	10.5	2.23	12.4	1.98	12.2	2.25
100030	3	3	17.9	2.61	19.5	3.71	13.7	2.07	18.1	2.58	13.3	2.35	12.8	3.46	9.71	1.79	12.9	2.3
100031	0	0	4.34	1.18	7.98	1.92	8.21	1.67	8.4	2.01	3.87	1.13	6.03	1.78	6.95	1.56	6.44	1.88
100032	3	3	70.5	5.01	73.3	6.32	56.7	3.99	62.9	4.34	79.8	5.25	80.6	6.48	65.3	4.27	71.4	4.59
100033	0	0	1.92	0.779	1.41	0.912	1.7	1.01	-2.2	1.09	1.7	0.74	4.35	1.28	5.35	1.42	4.69	1.75
100034	1	1	2.63	0.936	26.4	4.46	6.69	1.54	12.3	2.26	2.16	0.866	14.8	4.1	5.03	1.39	6.49	1.88
100035	0	0	0.65	0.47	1.29	1.26	2.3	1.09	1.57	1.49	0.932	0.545	1.7	1.3	2.19	1.08	0.553	1.39
100036	2	2	152	7.13	151	7.3	142	6.24	149	6.48	136	6.83	136	7	127	5.91	133	6.15
100037	3	3	7.04	1.66	7.71	3.63	8.53	1.69	8.55	2.02	7.45	1.69	10.9	3.75	9	1.73	11.1	2.19
100038	0	0	21.2	2.64	25.3	3.77	24.4	2.68	25.1	2.93	14.3	2.26	14.3	3.35	16.4	2.24	16.5	2.5
100039	2	2	13.4	2.19	14.9	3.92	12.9	2.01	15.3	2.43	12.9	2.16	17.3	4.01	11.8	1.94	16.9	2.52
100040	1	0	0	0	0	0	0	0	0	0	14.8	2.17	15.8	2.74	18.3	2.35	18.4	2.6
100041	1	0	0	0	0	0	0	0	0	0	23.9	2.74	26.4	3.37	28.5	2.88	28.8	3.09

**Note.** — This is a portion of the full table to provide form and guidance. 1. Object ID number in Table 7. 2. SExtractor flags in the 3.6  $\mu$ m image, 3. SExtractor flags in the 4.5  $\mu$ m image, 4. isophotal flux in the 3.6  $\mu$ m image, 5. error on isophotal flux, 6. total (Kron) flux in 3.6  $\mu$ m image, 7. error on total flux, 8. 3.6  $\mu$ m flux measured in 4''-diameter aperture, corrected to total, 9. error on 4''-diameter flux, 10. 3.6  $\mu$ m flux measured in 6''-diameter aperture, corrected to total, 11. error on 6''-diameter flux, 12. isophotal flux in the 4.5  $\mu$ m image, 13. error on isophotal flux, 14. total (Kron) flux in 4.5  $\mu$ m image, 15. error on total flux, 16. 4.5  $\mu$ m flux measured in 4''-diameter aperture, corrected to total, 17. error on 4''-diameter flux, 18. 4.5  $\mu$ m flux measured in 6''-diameter aperture, corrected to total, 19. error on 6''-diameter flux.

**Table 9**  
Photometry for SHELA Epoch 1 IRAC Catalogs

ID	Flags	Flags	$f_{\nu,ISO}^{(3.6)}$	$\sigma_{ISO}^{(3.6)}$	$f_{\nu,AUTO}^{(3.6)}$	$\sigma_{AUTO}^{(3.6)}$	$f_{\nu,4''}^{(3.6)}$	$\sigma_{4''}^{(3.6)}$	$f_{\nu,6''}^{(3.6)}$	$\sigma_{6''}^{(3.6)}$	$f_{\nu,ISO}^{(4.5)}$	$\sigma_{ISO}^{(4.5)}$	$f_{\nu,AUTO}^{(4.5)}$	$\sigma_{AUTO}^{(4.5)}$	$f_{\nu,4''}^{(4.5)}$	$\sigma_{4''}^{(4.5)}$	$f_{\nu,6''}^{(4.5)}$	$\sigma_{6''}^{(4.5)}$
(1)	(3.6 $\mu$ m)	(4.5 $\mu$ m)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	(14)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)
100020	3	2	0	0	0	0	0	0	0	0	506	13.6	514	13.8	385	10.3	463	11.4
100021	0	0	3.78	1.33	9.91	4.72	6.02	1.81	5.85	2.59	2.46	1.19	-2.32	nan	3.5	1.61	0.925	2.32
100022	2	2	8.11	1.94	9.44	4.99	10.6	2.13	13.4	2.96	4.43	1.67	10.2	5.01	5.82	1.8	5.77	2.59
100023	1	1	3.32	1.58	1.02	2.06	2.42	1.52	-0.875	2.22	5.19	1.73	6.8	2.41	5.91	1.81	5.83	2.59
100024	1	0	0	0	0	0	0	0	0	0	27.2	3.48	28.1	4.32	30.7	3.15	31.7	3.7
100025	0	0	2.4	1.18	3.13	3.91	3.14	1.58	3.39	2.46	3.04	1.25	5.76	4	5.29	1.76	4.44	2.52
100026	1	0	0	0	0	0	0	0	0	0	2.17	0.955	3.71	1.5	5.17	1.75	3.62	2.48
100027	0	0	22.4	3.24	26.4	4.47	22.5	2.78	25.4	3.47	24.2	3.32	26.3	4.46	26.6	2.98	27.5	3.55
100028	1	0	0	0	0	0	0	0	0	0	3.47	1.23	4.3	2.17	4.69	1.71	4.92	2.55
100029	1	0	0	0	0	0	0	0	0	0	9.98	2.07	13	3.1	13.4	2.3	13.9	2.98
100030	3	3	18.6	3.26	16.8	5.18	14.5	2.36	20.6	3.27	11.5	2.95	12.1	5.06	8.36	1.98	12.1	2.9
100031	0	0	4.28	1.35	7.92	2.57	8.11	1.96	8.37	2.72	3.75	1.29	5.89	2.46	6.7	1.86	6.27	2.62
100032	3	3	71.1	5.96	71.4	8.21	56.6	4.12	63.4	4.72	84.9	6.26	84.5	8.42	68.9	4.5	76	5.07
100033	1	0	0	0	0	0	0	0	0	0	0.851	0.695	2.37	1.4	2.18	1.5	0.543	2.3
100034	1	1	2.67	1.1	26	6.26	6.61	1.86	12.1	2.9	2.14	1.03	15.1	6.02	5.06	1.74	6.55	2.63
100035	0	0	0.233	0.435	-1.22	nan	0.9	1.38	-0.846	2.22	1.01	0.631	2.14	2.04	2.5	1.53	1.23	2.34
100036	2	2	148	8.09	149	8.45	141	6.3	148	6.72	147	8.08	146	8.4	128	6.01	137	6.5
100037	3	3	5.01	1.95	3.26	5.41	5.99	1.81	6.37	2.62	9.81	2.26	17.8	5.76	11.1	2.16	17.4	3.14
100038	0	0	18	2.9	19.1	4.97	20.9	2.7	21.1	3.29	13.7	2.69	14	4.83	17.2	2.51	15.4	3.05
100039	2	2	15	2.73	19.5	5.85	14	2.33	15.6	3.06	12.6	2.61	22.5	5.92	10.2	2.1	15.1	3.04
100040	1	0	0	0	0	0	0	0	0	0	16.3	2.54	17.9	3.73	20.4	2.68	20.3	3.26
100041	1	0	0	0	0	0	0	0	0	0	22	2.99	24.2	4.27	26.4	2.97	27	3.53

**Note.** — This is a portion of the full table to provide form and guidance. 1. Object ID number in Table 7, 2. SExtractor flags in the 3.6  $\mu$ m image, 3. SExtractor flags in the 4.5  $\mu$ m image, 4. isophotal flux in the 3.6  $\mu$ m image, 5. error on isophotal flux, 6. total (Kron) flux in 3.6  $\mu$ m image, 7. error on total flux, 8. 3.6  $\mu$ m flux measured in 4''-diameter aperture, corrected to total, 9. error on 4''-diameter flux, 10. 3.6  $\mu$ m flux measured in 6''-diameter aperture, corrected to total, 11. error on 6''-diameter flux, 12. isophotal flux in the 4.5  $\mu$ m image, 13. error on isophotal flux, 14. total (Kron) flux in 4.5  $\mu$ m image, 15. error on total flux, 16. 4.5  $\mu$ m flux measured in 4''-diameter aperture, corrected to total, 17. error on 4''-diameter flux, 18. 4.5  $\mu$ m flux measured in 6''-diameter aperture, corrected to total, 19. error on 6''-diameter flux.

**Table 10**  
Photometry for SHELA Epoch 2 IRAC Catalogs

ID	Flags	Flags	$f_{\nu,ISO}^{(3.6)}$	$\sigma_{ISO}^{(3.6)}$	$f_{\nu,AUTO}^{(3.6)}$	$\sigma_{AUTO}^{(3.6)}$	$f_{\nu,4''}^{(3.6)}$	$\sigma_{4''}^{(3.6)}$	$f_{\nu,6''}^{(3.6)}$	$\sigma_{6''}^{(3.6)}$	$f_{\nu,ISO}^{(4.5)}$	$\sigma_{ISO}^{(4.5)}$	$f_{\nu,AUTO}^{(4.5)}$	$\sigma_{AUTO}^{(4.5)}$	$f_{\nu,4''}^{(4.5)}$	$\sigma_{4''}^{(4.5)}$	$f_{\nu,6''}^{(4.5)}$	$\sigma_{6''}^{(4.5)}$
(1)	(2)	(3)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	(14)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)
100020	2	2	453	13	451	13.1	334	9.59	408	10.8	523	13.7	521	13.8	391	10.4	469	11.5
100021	0	0	1.05	1.02	11.7	4.88	3.25	1.62	6.56	2.71	4.23	1.38	7.81	4.77	7.25	1.92	6.32	2.69
100022	3	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100023	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100024	0	0	43.5	4.11	45.1	4.91	48.4	3.85	49.3	4.34	28.3	3.58	28.7	4.43	32.2	3.23	32.3	3.78
100025	0	0	4.27	1.38	1.29	3.97	6.68	1.88	4.18	2.58	4.44	1.4	9	4.23	6.81	1.89	6.67	2.71
100026	0	1	2.09	0.941	3.5	1.5	4.95	1.76	4.36	2.59	0	0	0	0	0	0	0	0
100027	0	0	16.8	3.06	15.8	4.24	17.9	2.57	17.8	3.22	22	3.28	22.8	4.46	25.6	2.94	24.8	3.5
100028	0	1	1.7	1.02	2.02	2.09	3.8	1.66	3.82	2.56	0	0	0	0	0	0	0	0
100029	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100030	3	3	17.3	3.28	19.1	5.34	13.8	2.34	16.1	3.15	18.5	3.33	20	5.36	13.2	2.31	18.4	3.24
100031	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100032	3	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100033	0	1	1.9	0.873	1.42	1.33	1.72	1.49	-2.22	2.22	0	0	0	0	0	0	0	0
100034	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100035	0	1	0.744	0.567	-0.221	nan	1.39	1.46	-0.559	2.32	0	0	0	0	0	0	0	0
100036	2	2	146	8.11	145	8.43	139	6.27	146	6.71	125	7.76	124	8.09	126	5.98	129	6.35
100037	3	3	7.01	2.12	7.93	5.6	8.07	1.98	7.97	2.77	5.79	2.04	8.11	5.61	7.35	1.93	8.29	2.79
100038	0	1	22.7	3.15	30.6	5.38	26.6	2.99	27.2	3.59	0	0	0	0	0	0	0	0
100039	3	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100040	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100041	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

**Note.** — This is a portion of the full table to provide form and guidance. 1. Object ID number in Table 7, 2. SExtractor flags in the 3.6  $\mu$ m image, 3. SExtractor flags in the 4.5  $\mu$ m image, 4. isophotal flux in the 3.6  $\mu$ m image, 5. error on isophotal flux, 6. total (Kron) flux in 3.6  $\mu$ m image, 7. error on total flux, 8. 3.6  $\mu$ m flux measured in 4''-diameter aperture, corrected to total, 9. error on 4''-diameter flux, 10. 3.6  $\mu$ m flux measured in 6''-diameter aperture, corrected to total, 11. error on 6''-diameter flux, 12. isophotal flux in the 4.5  $\mu$ m image, 13. error on isophotal flux, 14. total (Kron) flux in 4.5  $\mu$ m image, 15. error on total flux, 16. 4.5  $\mu$ m flux measured in 4''-diameter aperture, corrected to total, 17. error on 4''-diameter flux, 18. 4.5  $\mu$ m flux measured in 6''-diameter aperture, corrected to total, 19. error on 6''-diameter flux.



**Table 11**  
Photometry from SHELA Epoch 3 IRAC Catalogs

ID	Flags	Flags	$f_{\nu,ISO}^{(3.6)}$	$\sigma_{ISO}^{(3.6)}$	$f_{\nu,AUTO}^{(3.6)}$	$\sigma_{AUTO}^{(3.6)}$	$f_{\nu,4''}^{(3.6)}$	$\sigma_{4''}^{(3.6)}$	$f_{\nu,6''}^{(3.6)}$	$\sigma_{6''}^{(3.6)}$	$f_{\nu,ISO}^{(4.5)}$	$\sigma_{ISO}^{(4.5)}$	$f_{\nu,AUTO}^{(4.5)}$	$\sigma_{AUTO}^{(4.5)}$	$f_{\nu,4''}^{(4.5)}$	$\sigma_{4''}^{(4.5)}$	$f_{\nu,6''}^{(4.5)}$	$\sigma_{6''}^{(4.5)}$
(1)	(3.6 $\mu$ m)	(4.5 $\mu$ m)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	(14)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)	( $\mu$ Jy)
100020	3	2	0	0	0	0	0	0	0	0	529	13.8	528	13.9	395	10.4	474	11.6
100021	0	0	2.94	1.25	7.83	4.69	3.94	1.66	4.06	2.53	1.06	1.02	-0.863	nan	2.14	1.51	2.93	2.47
100022	2	2	2.87	1.55	3.54	4.86	3.65	1.64	3.73	2.52	5.37	1.76	6.02	4.92	7.6	1.94	7.79	2.73
100023	1	1	4	1.65	4.04	2.28	3.03	1.59	4.67	2.57	5.94	1.8	7.72	2.48	7.25	1.91	8.22	2.75
100024	1	0	0	0	0	0	0	0	0	0	24.4	3.4	23.2	4.2	31.4	3.19	28	3.59
100025	0	0	3.38	1.29	15.7	4.36	6.51	1.86	11.6	2.91	1.01	1.01	-3.61	nan	0.423	1.35	-2.04	2.19
100026	1	0	0	0	0	0	0	0	0	0	2.67	1.02	3.52	1.49	4.62	1.72	4.44	2.55
100027	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100028	1	0	0	0	0	0	0	0	0	0	3.16	1.19	3.74	2.17	4.47	1.71	3.16	2.49
100029	1	0	0	0	0	0	0	0	0	0	7.98	1.95	6.38	2.83	10.7	2.15	9.06	2.79
100030	3	3	16.4	3.2	21.2	5.32	12.3	2.24	16.4	3.12	9.7	2.9	4.07	4.86	7.64	1.94	7.77	2.73
100031	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100032	3	3	68.3	5.92	74.9	8.2	56.2	4.11	60.7	4.66	73.5	6.04	75	8.2	61.5	4.28	66.2	4.82
100033	1	0	0	0	0	0	0	0	0	0	2.55	0.97	6.13	1.73	8.2	1.98	8.53	2.76
100034	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100035	0	0	0.787	0.58	6.71	2.35	5.44	1.78	8.56	2.76	0.86	0.596	1	2	1.78	1.48	-0.328	2.29
100036	2	2	159	8.29	158	8.6	145	6.4	151	6.8	135	7.88	135	8.23	127	6.01	133	6.43
100037	3	3	8.98	2.23	11.1	5.61	11.4	2.19	11.2	2.89	7.24	2.12	7.34	5.52	8.96	2.03	8.64	2.77
100038	0	0	23.4	3.16	26.3	5.19	26.1	2.96	27.7	3.58	14.6	2.76	14	4.86	15.4	2.42	17.1	3.15
100039	2	2	12.2	2.61	10.4	5.64	12.1	2.23	15.1	3.07	12.6	2.63	10.2	5.64	12.9	2.28	17.8	3.18
100040	1	0	0	0	0	0	0	0	0	0	11.9	2.31	11.7	3.55	14.4	2.37	14.4	3.04
100041	1	0	0	0	0	0	0	0	0	0	25.7	3.17	27.7	4.41	30.5	3.15	30.2	3.67

**Note.** — This is a portion of the full table to provide form and guidance. 1. Object ID number in Table 7, 2. SExtractor flags in the 3.6  $\mu$ m image, 3. SExtractor flags in the 4.5  $\mu$ m image, 4. isophotal flux in the 3.6  $\mu$ m image, 5. error on isophotal flux, 6. total (Kron) flux in 3.6  $\mu$ m image, 7. error on total flux, 8. 3.6  $\mu$ m flux measured in 4''-diameter aperture, corrected to total, 9. error on 4''-diameter flux, 10. 3.6  $\mu$ m flux measured in 6''-diameter aperture, corrected to total, 11. error on 6''-diameter flux, 12. isophotal flux in the 4.5  $\mu$ m image, 13. error on isophotal flux, 14. total (Kron) flux in 4.5  $\mu$ m image, 15. error on total flux, 16. 4.5  $\mu$ m flux measured in 4''-diameter aperture, corrected to total, 17. error on 4''-diameter flux, 18. 4.5  $\mu$ m flux measured in 6''-diameter aperture, corrected to total, 19. error on 6''-diameter flux.

**Table 13**  
SDSS Stripe 82 Coadd Photometry for sources matched to SHELA

ID	SDSS ID	SDSS RA (deg)	SDSS DEC (deg)	TYPE	SDSS FLAGS	$u$ (mag)	$\sigma_u$ (mag)	$g$ (mag)	$\sigma_g$ (mag)	$r$ (mag)	$\sigma_r$ (mag)	$i$ (mag)	$\sigma_i$ (mag)	$z$ (mag)	$\sigma_z$ (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)
100020	8647474690342256787	17.701219	-1.132932	3	103347650576	19.377	0.016	18.439	0.004	17.740	0.003	17.434	0.003	17.175	0.006
100024	8647474690341142937	15.221294	-1.125124	6	34628174080	24.438	0.551	21.697	0.025	20.347	0.009	19.601	0.007	19.192	0.014
100029	8647474690343175313	19.864137	-1.127867	3	2450547277824	22.385	0.151	22.097	0.066	21.192	0.029	20.948	0.037	20.984	0.123
100036	8647474690342650333	18.621063	-1.131875	3	103347650560	22.802	0.243	21.048	0.027	19.731	0.009	19.213	0.010	18.837	0.019
100038	8647474690342651302	18.717668	-1.128721	3	70439879574360	25.901	3.217	24.574	0.547	23.981	0.294	23.269	0.231	22.215	0.290
100041	8647474690343175190	19.928157	-1.128586	3	68987912448	27.804	3.907	23.321	0.116	21.927	0.031	21.062	0.023	20.654	0.051
100044	8647474690342258061	17.794618	-1.127763	3	68987912448	23.361	0.328	22.871	0.114	21.835	0.042	21.323	0.039	21.003	0.095
100048	8647474690342782172	18.936888	-1.128068	3	281543964622848	24.181	0.547	23.570	0.170	23.485	0.150	23.571	0.253	24.049	1.245
100049	8647474690342717098	18.867537	-1.128079	3	68987912448	26.888	3.737	23.853	0.183	23.418	0.112	22.700	0.091	22.371	0.220
100052	8647474690341797917	16.668147	-1.135992	3	34628173840	16.636	0.003	15.029	0.003	14.560	0.003	14.510	0.003	14.477	0.003

**Note.** — This is a portion of the full table to provide form and guidance. 1. Object ID number in Table 7, 2. SDSS ID number, 3. SDSS right ascension (J2000) in decimal degrees, 4. SDSS declination (J2000) in decimal degrees, 5. SDSS Type (common values are Type=3 for galaxy and Type=6 for star), 6. SDSS Flags value, 7. SDSS  $u$  magnitude, 8. error on  $u$  magnitude, 9. SDSS  $g$  magnitude, 10. error on  $g$  magnitude, 11. SDSS  $r$  magnitude, 12. error on  $r$  magnitude, 13. SDSS  $i$  magnitude, 14. error on  $i$  magnitude, 15. SDSS  $z$  magnitude, 16. error on  $z$  magnitude