

# Star Formation

## Answering Fundamental Questions During the Spitzer Warm Mission Phase

Lori Allen CfA  
John Carpenter, Caltech  
Lee Hartmann, University of Michigan  
Michael Liu, University of Hawaii  
Tom Megeath, University of Toledo  
Luisa Rebull, Spitzer Science Center  
John Stauffer, Spitzer Science Center  
Steve Strom NOAO

## Spitzer Key Contributions to Date

- Provide a census of forming stars in nearby molecular clouds
  - Sensitivity to  $\sim 10 M_{\text{Jupiter}}$  for star-forming regions to distances  $d \sim 500$  pc
  - Includes protostars and young stars surrounded by circumstellar disks at various evolutionary phases




## Spitzer Key Contributions to Date

- Provide basis for addressing key problems
  - How are initial core conditions related to outcome stellar properties?
  - What is the range of lifetimes available for planet-building?
  - Is there evidence for ongoing planet formation in disks?

## Spitzer Key Contributions to Date

- Provide the basis for answering
  - What conditions in molecular clouds lead to the observed population mix?
  - How does the cloud environment influence star formation and disk lifetimes?
- Provide a census of stars surrounded by circumstellar accretion disks
  - Span ages from  $t < 1$  Myr to  $t \sim 25$  Myr



## Spitzer Key Contributions to Date

- Provide direct evidence of accretion disk evolution
  - Grain settling to the midplane
  - Grain growth
  - Evolution in the distribution of solid material in the inner disk
- Provide the basis for answering
  - Timescale(s) available for planet-building
  - Timescales for key physical and chemical processes in disks
  - When, where and under what conditions planet-building may take place



## Focus of Spitzer Observations to Date

- Deep observations of star-forming regions directly associated with well-known star-bearing molecular clouds located within 500 pc of the sun



## Focus of Spitzer Observations to Date

- Targeted observations of more distant regions located at 1-2 kpc
  - Generally focused on known centers of star formation or dense molecular gas



## What Spitzer has yet to do

- Explore larger regions surrounding the active star-forming centers in nearby clouds
  - Environs of T- and OB- associations in which multiple episodes of star formation have taken place over timescales of 1-20 Myr
- Explore giant star-forming complexes found in the Carina and Perseus arms
  - Are sites of star-forming activity analogous to those in other galaxies
  - Provide evidence of the processes that initiate and propagate star formation
- Explore star-formation on the scale of a spiral arm
  - Understand how the stellar populations comprising galactic disks arise
- Explore variable sources at a variety of cadences and over long timescales
  - Gain insights into the fundamental physics of the accretion/stellar assembly process

## The Power of Spitzer During the Warm Mission Phase

- Sensitivity sufficient to detect excess IR emission arising from protostars and young stars still surrounded by circumstellar accretion disks
  - Down to or below the HBL in nearby ( $d < 500$  pc) OB- and T- associations
  - Down to  $\sim 0.3 M_{\text{sun}}$  in more distant ( $d \sim 2$  kpc) clouds
- Ability to map several hundred square degrees in of order a thousand hours
- Angular resolution sufficient to
  - Resolve all but the densest regions of forming groups and clusters
  - Identify contaminating background galaxies whose colors mimic those of young stars

## Spitzer Warm Mission: Three Proposed Programs

- Survey of Selected OB Associations
- Survey of Distant GMCs & a representative region of a spiral arm
- A Search for Variability Patterns among YSOs


## Spitzer Warm Mission: Survey of OB Associations

- Goals
  - Understand the number and spatial distribution of star-forming episodes
  - Understand the link of these patterns to initial conditions in molecular clouds
  - Determine the timescales for circumstellar disk evolution
    - Sample will include stars spanning a wide range of ages (up to 20 Myr) and masses


## Spitzer Warm Mission: Survey of OB Associations

- Approach
  - Map areas of 50-200 square degrees of 3 representative OB associations
  - Survey depth: 12  $\mu$ Jy (IRAC 1) and 18  $\mu$ Jy (IRAC 2)
  - Will detect stars surrounded by accretion disks to masses  $\sim 0.08$  Myr at ages  $t \sim 5$  Myr
- Motivation for a Warm Mission Survey
  - Thus far, only selected regions (largely confined to dense molecular gas) have been mapped
  - Older star-forming episodes have largely been excluded
  - We lack information re the patterns of star-formation and timescales associated with each episode
  - Constraints on disk lifetimes as a function of mass and age are not yet robust
    - Samples are heavily biased toward regions younger than 5 Myr

## Spitzer Warm Mission: Survey of OB Associations



## Spitzer Warm Mission: Survey of OB Associations




# Spitzer Warm Mission: Survey of OB Associations

- Expected results
  - Complete surveys of protostars and stars surrounded by circumstellar accretion disks
  - Morphology will provide guide to sequence and possible triggers of star-forming events
  - Essential constraints on accretion disk evolutionary, and thus planet-building timescales
- Complementary ground-based observations motivated by Spitzer
  - Multi-epoch JHK imaging surveys
    - Detect young stars that lack disks via variability
    - Complementary tool for weeding out galaxies from candidate stars surrounded by disks or envelopes
  - High resolution spectroscopy
    - Kinematics of stellar population (key link to where and how stars formed and dispersed)
    - Accretion rate vs time and stellar mass
    - Infall rates in protostellar envelopes
  - ALMA observations of disk masses (dust continuum) + envelope properties

# Spitzer Warm Mission: Survey of OB Associations


Value of Spitzer: Power to Identify Young Stars from IRAC colors





## Spitzer Warm Mission: Survey of OB Associations

Value of Spitzer: Power to Identify Young Stars from IRAC colors



## Spitzer Warm Mission: Survey of GMCs + Outer Galaxy Arm

- Goals
  - Examine star-formation in regions more analogous to those in other galaxies
    - Include clusters of richness and density far exceeding that of the Orion Nebula Cluster
  - Understand the pattern and timescales for star-forming events within GMCs
  - Understand the relationship between forming stars, propagating HII regions and superbubbles
  - Examine star-forming patterns on the scale of a spiral arm
    - Understand the extent of star-formation outside the boundaries of dense GMCs

## Spitzer Warm Mission: Survey of GMCs + Outer Galaxy Arm


- Approach
  - Map area of ~300 square degrees of 5 representative GMCs (in Perseus and Carina arms)
  - Map area of ~300 square degrees in a relatively unconfused region of the Perseus arm
    - Chosen solely to lie within a fixed  $A_V$  contour; no bias toward known star-forming regions
  - Survey depth: 12  $\mu\text{Jy}$  (IRAC 1) and 18  $\mu\text{Jy}$  (IRAC 2)
  - Will detect stars surrounded by accretion disks to masses  $\sim 0.3$  Myr at ages  $t \sim 5$  Myr
- Motivation
  - To date, selected regions of GMCs have been mapped
    - focused on dense molecular gas and known signposts of star-formation
  - Large complexes analogous to those populating spiral arms in actively star-forming galaxies have not yet been mapped completely
  - Objective studies of star-formation on the scale of a spiral arm have been minimal

## Spitzer Warm Mission: Survey of GMCs + Outer Galaxy Arm



Multicolor IRAC Map of W5 (1.5 x 2°)

## Spitzer Warm Mission: Survey of GMCs + Outer Galaxy Arm



## Spitzer Warm Mission: Survey of GMCs + Outer Galaxy Arm

- Complementary ground- and space- based observations motivated by Spitzer
  - ALMA, CARMA, SMA and SCUBA-2 observations of
    - molecular cloud physical, chemical and kinematic properties (understand initial conditions)
    - Individual star-forming cores (understand relationship between core & outcome stellar properties)
  - High resolution near- and mid-IR spectroscopy
    - Determine conditions in the inner regions of star-forming cores
    - Determine stellar kinematics and relate to molecular cloud properties
  - Chandra, XMM and near-IR ground-based imaging surveys
    - Complete population survey in selected regions by locating young stars that have lost their disks
  - Moderate resolution ground-based spectroscopy (optical and near-IR)
    - Classification spectra needed to place stars in HRD: determine masses and ages
    - Determine sequence of star-formation
  - JWST imaging and spectroscopy in dense forming clusters
    - Determine stellar properties (e.g. IMF) in regions similar to dense clusters in other galaxies


## Spitzer Warm Mission: Variability Patterns among YSOs

- Goals:
  - Determine variability patterns among stars surrounded by protostellar cores and/or accretion disks
  - Diagnose source of variability
    - Warm or cool star spots
    - Variable obscuration by envelope/disk material
    - Variability driven by changes in accretion rate and heating in the inner disk
    - Variability driven by envelope/disk interactions

## Spitzer Warm Mission: Variability Patterns among YSOs

- Approach (example program):
  - Monitor a 1x3 degree region centered on the Trapezium
  - Region can be mapped on four hour timescale
    - 400 non-overlapping IRAC frames
    - 12 sec integration HDR mode (yields  $50\sigma$  at  $4.5\mu\text{m}$ ): sufficient to reach stars at the HBL
  - Repeat observations (a) six times per day for one 44 day visibility window of Orion, and then (b) repeat once per day for the next two visibility windows.
  - Total program time: ~ 1300 hours
- Motivation for a Spitzer Warm Mission Survey
  - 2MASS survey reveals variability among 100s of stars on timescales from hours to months
  - Spitzer has a unique combination of sensitivity, areal coverage & photometric precision
    - Ground-based L-band surveys map much smaller areas; precision limited to ~ 10%

## Spitzer Warm Mission: Variability Patterns among YSOs



## Spitzer Warm Mission: Variability Patterns among YSOs

- Possible results:
  - Understand factors that control variability in the inner (0.1-1 AU) disk regions
  - Understand variability during the infall phase
  - Understand the possible role of events driven by orbiting planets within accretion disk
- Complementary ground-based observations motivated by Spitzer
  - JHK imaging surveys aimed at sorting out stellar from disk-driven events
  - High resolution spectroscopy of selected objects to monitor emission line profile variability during same windows as IRAC photometry

## Spitzer Warm Mission: Comparison to Other Facilities

- Conclusions
  - Large area surveys with JWST impractical
    - Small FOV and downlink data volume constraints
    - Reserve JWST for crowded/selected regions
  - Surveys from Keck cannot reach desired sensitivities in practical times
    - Keck AO may be important for selected sources
  - WISE lacks sensitivity to survey below HBL in nearby regions
  - WISE lacks angular resolution needed in more distant regions
  - Spitzer Warm hits a ‘sweet spot’ in sensitivity and angular resolution

## Spitzer Warm Mission: Comparison to Other Facilities

